



# Enhancing Crop Saline Stress Tolerance and Soil Remediation with Phytobeneficial Bacteria: Diverse Approaches for Improvement

Divyashree <sup>a\*</sup>, A. Deepasree <sup>b</sup>, T. L. Shivananda <sup>c</sup>  
and Fidha Husna K <sup>d</sup>

<sup>a</sup> Department of Plant Pathology, NMCA, Navsari Agricultural University, Navsari, Gujarat-396450, India.

<sup>b</sup> Department of Soil Science and Agricultural Chemistry, NMCA, Navsari Agricultural University, Navsari, Gujarat-396450, India.

<sup>c</sup> Department of Molecular Biology and Agricultural Biotechnology, University of Agricultural science, Raichur, Karnataka- 584104, India.

<sup>d</sup> Department of Soil Science and Agricultural Chemistry, Kerala Agricultural University, Kerala-680656, India.

## Authors' contributions

This work was carried out in collaboration among all authors. Author Divyashree designed the study, performed the statistical analysis, wrote the protocol, and first draft of the manuscript. Authors AD and TLS managed the analyses of the study. Author FHK managed the literature searches. All authors read and approved the final manuscript.

## Article Information

DOI: <https://doi.org/10.9734/ajsspn/2024/v10i2312>

## Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/114466>

Review Article

Received: 14/01/2024  
Accepted: 17/03/2024  
Published: 25/05/2024

\*Corresponding author: Email: [divyasiriii7@gmail.com](mailto:divyasiriii7@gmail.com);

**Cite as:** Divyashree, Deepasree, A., Shivananda, T. L., & Husna K, F. (2024). Enhancing Crop Saline Stress Tolerance and Soil Remediation with Phytobeneficial Bacteria: Diverse Approaches for Improvement. *Asian Journal of Soil Science and Plant Nutrition*, 10(2), 539–558. <https://doi.org/10.9734/ajsspn/2024/v10i2312>

## ABSTRACT

Soil salinity, a pervasive issue exacerbated by factors like irrigation and climate change, poses a significant threat to global food security. The accumulation of salts not only hampers crop growth and yield but also jeopardizes the livelihoods of millions who depend on agriculture for sustenance. Elevated salt levels in saline soils induce osmotic, ionic, oxidative, and water stress in plants. Implementing biological solutions offers the most dependable and sustainable method to safeguard food security while reducing reliance on agrochemicals which hampers various physiological and metabolic processes in plants. To ensure optimal plant growth under such changing conditions, implementing biological solutions (Rhizobacteria) offers the most dependable and sustainable method to safeguard food security while reducing reliance on agrochemicals must be integrated into agricultural practices. This chapter concisely explores the mechanisms and utilization of beneficial microorganisms in both plants and soil to mitigate salt stress. It also addresses the current limitations and suggests potential areas for improvement in future research.

*Keywords: Saline soil; rhizobacteria; remediation; Pseudomonas spp.*

## 1. INTRODUCTION

The population of Earth reached 8.1 billion people in 2010; if growth continues at its current rate, that number is predicted to reach 9.7 billion people by 2050 (Projections of population growth). Additionally, water, air and soil pollution are responsible for about 40% of deaths globally and environmental deterioration like this, together with population growth, are thought to be important factors in the rapid rise in human disease worldwide. Various abiotic factors such as temperature, salinity, drought, pesticide and fertilizer usage, soil pH, and heavy metal contamination can impede crop productivity [1-3]. Out of all of these, soil salinity's worldwide effect on crop productivity has emerged as a major barrier. Human activity has increased the development of soil salinization during the last few decades [4-6]. Important soil activities like respiration, residue breakdown, nitrification, denitrification, soil biodiversity, and microbial activity are all impacted by the extreme soil salinization [7]. "Increased soil salinity and decreased crop output are also observed in areas with excessive fertiliser application" [8]. "The technique of removing salt from saline soil is labor-intensive and expensive" [9]. "For quite some time, the rehabilitation of saline soils has primarily relied on physical and chemical techniques. Within the realm of physical processes, soluble salts within the root zone are extracted through methods such as scraping, flushing, and leaching" [10]. "Nevertheless, chemical methods often involve the utilization of gypsum and lime as neutralizing agents to mitigate saline soil conditions" [11]. "However, these methods are deemed unsustainable and are considered inefficient,

particularly when the salt concentration reaches excessively high levels, The common practice of cultivating salt-tolerant crop varieties, such as barley and canola, on saline soils is widespread" [12]. "Nevertheless, due to their limited salt tolerance profile, these crops have a restricted global distribution and cannot be effectively utilized in soils with moderate to high electrical conductivity (EC) levels also highlighted that despite vigorous efforts from the research community, only few salt tolerance genes have been identified having real applications in improving productivity of saline soils" [13].

"Therefore, achieving viable crop yields in saline soils is imperative. In addition to utilizing salt-tolerant varieties or chemical neutralization methods, it's essential to incorporate sustainable approaches. In the last few years, research showed that the use of salt-tolerant plant growth promoting rhizobacteria (ST-PGPR) and halotolerant rhizobacteria (HT-rhizobacteria) in saline agriculture can be harnessed for enhancing productivity and improving soil fertility as well". As they significantly impact biogeochemical cycles, soil fertility, and plant health, they play a crucial role in influencing plant growth and the uptake of nutrients. This review critically examines the role of salt-tolerant plant growth-promoting rhizobacteria (ST-PGPR) and halo tolerant rhizobacteria in responding to salt-affected soil and their beneficial effects on key crops. It delves into their mechanisms for remediating salt-affected soil under diverse environmental conditions. The present review focuses on the enhancement of productivity under stressed conditions and increased resistance of plants against salinity stress by

application of plant growth promoting microorganisms.

The utilization of Plant Growth-Promoting Rhizobacteria (PGPR) has been expanded to remediate contaminated soils in conjunction with plants. Therefore, there is a pressing necessity to augment the effectiveness of limited external inputs by optimizing the combinations of beneficial bacteria within sustainable agricultural production systems. This review delves into the significance of soil-beneficial bacteria and their contributions to promoting plant growth through both direct and indirect mechanisms. A deeper understanding of these varied mechanisms will contribute to establishing these bacteria as invaluable allies in the future of agriculture.

## 2. MECHANISM OF CROP SALINE STRESS TOLERANCE BY SOIL BENEFICIAL BACTERIA

Soil salinity poses a significant challenge for irrigated agriculture. In hot and arid regions across the globe, soils often exhibit high salinity levels, resulting in limited agricultural productivity. It's worth noting that all soils inherently contain some amount of water-soluble salts [14]. Soluble salts are a form of essential nutrients that plants absorb; nevertheless, an overabundance of them can seriously impede plant growth. Global natural resources have suffered greatly as a result of land degradation processes over the past century, whether they are physical, chemical, or biological. Compacted soil, contamination from both organic and inorganic sources, and a decrease in microbial variety and activity are a few of these problems [15-17]. "Salinity destructively interrupts the physical and chemical properties of soil as well as affects crop growth to a higher extent" [18].

To address this issue, beneficial microorganisms called plant growth-promoting rhizobacteria (PGPR) could serve a vital function. These rhizospheric bacteria have the ability to efficiently colonize plant roots, thereby contributing to soil fertility maintenance. They provide a promising alternative to traditional inorganic fertilizers and pesticides [19]. Previous reports have highlighted the efficacy of PGPR in enhancing the growth of different crops under conditions of salt stress [20,21]. The initial selection of locally-isolated salt-tolerant PGPR for addressing salinity is essential to guarantee their effectiveness. Studies have shown that indigenous strains are more proficient in enhancing plant resistance to

salinity stress compared to PGPR from non-saline ecosystems [22]. These beneficial microbes employ various mechanisms to mitigate salt stress, such as regulating the Na<sup>+</sup>/K<sup>+</sup> ratio by secreting extracellular polymeric substances known as exopolysaccharides (EPS), this mechanism enhances their survival in unfavorable soil conditions [23,24].

"The previous findings have reported that several bacterial genera, including *Pseudomonas*, *Bacillus*, *Burkholderia*, *Enterobacter*, *Microbacterium*, *Planococcus*, *Halomonas* could produce EPS (Exopolysaccharides) in salt stress condition" [25,26]. The exopolysaccharides play a vital role in bacterial aggregation or flocculation, chelates the various cations including Na<sup>+</sup> [27,28] facilitating the production of yield, this process involves the specific adsorption of the polymeric segment and polymer bridging between cells [29,30]. Additionally, EPS are highly beneficial in the formation of bacterial biofilms and enhancing bacterial colonization on plant root surfaces [31]. "Exopolysaccharides are able to lessen the hostile effect of osmotic-stress by augmenting fresh weight, dry weight and water content in plants were analysed statically" [32]. "In addition to that, PGPR are able to produce multiple plant growth-promoting properties such as indole acetic acid production, biological nitrogen fixation, solubilization of soil phosphorus (P) and potassium (K), and production of siderophores and hydrolyzing enzymes under salt stress condition" [33-36] "Plants treated with Exo-poly saccharides (EPS) producing bacteria display increased resistance to water and salinity stress due to improved soil structure" [37]. EPS can also bind to cations including Na<sup>+</sup> thus making it unavailable to plants under saline conditions.

"The SEM observations supported all these salt-tolerance attributes, revealing the bacterial capacity to produce EPS, facilitate flocculation, and form biofilms when subjected to saline conditions compared to non-saline environments. Bacterial cells were observed to associate with the plant root system, notably enhancing moisture retention capacity and bolstering the defense system against various abiotic stresses. Previous research also noted a reduction in bacterial EPS and biofilm formation with increased NaCl concentration" [38]. "The detrimental effects of salinity can be mitigated through the application of salt-tolerant PGPR, as demonstrated in this greenhouse trial. This intervention notably enhanced the

photosynthesis of all three rice varieties, resulting in increased grain yield under saline conditions” [39,40] “Soils experiencing salt stress are recognized for their ability to inhibit plant growth”. “In their natural habitat, plants are colonized by both endocellular and intracellular microorganisms” [41]. “The rhizosphere microorganisms, especially beneficial bacteria and fungi, have the potential to enhance plant performance in stressful environments, thereby directly and indirectly improving yields, Certain PGPR can directly stimulate plant growth and development by supplying fixed nitrogen, phytohormones, iron sequestered by bacterial siderophores, and soluble phosphate” [42]. “Others indirectly benefit plants by protecting them against soil-borne diseases, primarily caused by pathogenic fungi” [43]. “Soil salinization presents a significant challenge to agricultural productivity worldwide. Crops cultivated in saline soils face issues such as high osmotic stress, nutritional imbalances and toxicities, poor soil structure, and decreased crop yields. Studies has confirmed that salt-tolerant plant growth promoting rhizobacteria (ST-PGPR) are capable of producing various phytohormones, including auxins, gibberellins, and cytokinins” [44]. Additionally, they synthesize ACC deaminase [45], secondary compounds such as exopolysaccharides [46,47]. and osmolytes (proline, trehalose, and glycine betaines) [48,49]. “Furthermore, these bacteria play a role in regulating plant defense systems and activating the plant's antioxidative enzymes under salt stress” [50,51].

Numerous studies have highlighted potential mechanisms of salt tolerance in PGPR, particularly when they function as endophytes. However, it's important to note that only a small portion of these beneficial bacteria are able to penetrate the root cell, and the interaction between plants and microbes primarily takes place within the 5mm rhizosphere zone [52]. Salt-tolerant bacterial strains have demonstrated elevated nitrogenase activity in saline environments and possess the capability to synthesize osmolytes. These osmolytes help maintain cell turgidity and support metabolism in adverse conditions [53,54].

Salinity-induced nutritional imbalance poses a challenge to plant growth and productivity, particularly affecting phosphorus (P) uptake and transport. To address P deficiency in salt-affected soils, P fertilizers are commonly recommended. However, employing salt-tolerant

P-solubilizing rhizobacteria can significantly enhance P availability in saline soils [55]. In recent studies, plant growth-promoting (PGP) bacteria have been discovered to boost plant tolerance to salinity, particularly those bacteria that are associated with plants. Endophytic actinomycetes are particularly intriguing due to their dual role. Not only do they produce various bioactive secondary metabolites, safeguarding plants against infectious diseases, but they also exhibit the capacity to enhance plant growth. They carry several plant growth-promoting (PGP) traits, including the production of siderophores for iron acquisition, synthesis of plant hormones like auxins and cytokinins, and the solubilization of phosphate and other minerals to provide nutrients [56,57]. Furthermore, they assist in plant growth under stressful conditions induced by drought, heavy metals, flooding and high salinity by alleviating stress associated with ethylene through the production of 1-aminocyclopropane-1-carboxylate (ACC) deaminase. [58] correlated the role of proline accumulation with drought and salt tolerance in plants, Through these studies, we gain insight into the role of released hormones, enzymes, and other metabolic substances in alleviating salt stress conditions with the assistance of numerous rhizobacteria.

### **3. BACTERIAL MICROBES: SALT STRESS ALLEVIATION TOOL IN IMPORTANT CROPS**

“Several strategies have been developed in order to decrease the toxic effects caused by high salinity on plant growth, including plant genetic engineering, and recently the use of plant growth-promoting bacteria (PGPB)” [59].“The role of microorganisms in plant growth promotion, nutrient management and disease control is well known and well established. These beneficial microorganisms colonize the rhizosphere/endorhizosphere of plants and promote growth of the plants through various direct and indirect mechanisms” [60,61]. Previous studies suggest that utilization of PGPB has become a promising alternative to alleviate plant stress caused by salinity [62] and the role of microbes in the management of biotic and abiotic stresses is gaining importance.

#### **3.1 Cereal Crops**

Cereal crops serve as the primary sources of energy and protein in the human diet, cultivated

in significantly larger quantities worldwide compared to other crops. Major cereal crops include wheat, maize, rice, barley, oats, sorghum, and millet. Despite their importance, only a few of these crops exhibit salt tolerance. Traditional methods such as conventional breeding, marker-assisted selection, and genetic engineering have been successful in enhancing yields in saline soils, but primarily for wheat and rice over the past decades. It has been observed that the application of Salt-Tolerant Plant Growth-Promoting Rhizobacteria (ST-PGPR) in saline soil not only aids in crop survival but also enhances yields across a diverse array of cereal crops [63,64].

Similarly, the combination of *Pseudomonas pseudoalcaligenes*, an endophytic bacterium, with *Bacillus pumilus* in the rhizosphere of paddy plants was more effective in protecting the plants from abiotic stress during early growth stages. This combination induced the production of osmoprotectant and antioxidant proteins, surpassing the effects of either rhizospheric or endophytic bacteria alone. The plants inoculated with the endophytic bacterium *P. pseudoalcaligenes* exhibited notably elevated levels of glycine betaine-like quaternary compounds and increased shoot biomass, particularly evident at lower salinity levels [51]. This study demonstrates that the detrimental effects of salinity stress can be mitigated through the application of salt-tolerant Plant Growth-Promoting Rhizobacteria (PGPR). In a glasshouse trial, this approach significantly enhanced the photosynthetic activity of all three rice varieties, resulting in higher grain yields under saline conditions [39]. The inoculation of endophytic *Streptomyces* sp. GMKU 336, which produces 1-aminocyclopropane-1-carboxylate deaminase (ACCD), into rice plants leads to improved growth and enhanced salt tolerance. This is accomplished by using ACCD activity to lower ethylene levels, which aids in the plants' ability to scavenge reactive oxygen species (ROS), maintain a balanced ion content, and control osmotic pressure [65]. The research conducted by Ji, et al. [66]. This highlights the significant role of *Glutamicibacter* spp. YD01 in mitigating the detrimental effects of salt stress on the growth and development of rice plants. This is achieved through the regulation of phytohormone (ethylene) levels and the accumulation of reactive oxygen species (ROS), as well as maintaining ion balance, enhancing photosynthetic capacity, and promoting the expression of stress-responsive genes.

Similarly, the inoculation of endophytic *Methylobacterium oryzae* CBMB20 into salt-stressed rice plants, as observed in the study by Chatterjee et al. (2019), enhances photosynthesis and reduces emissions of stress-related volatiles. This is attributed to the modulation of ethylene-dependent responses and the activation of vacuolar H<sup>+</sup>-ATPase. Certain strains of *Rhodopseudomonas palustris*, such as TN114, show promise in facilitating the easier and more affordable growth of rice in saline soil conditions. This is attributed to the presence of 5-aminolevulinic acid (ALA) in the examined supernatants, which has a positive effect on rice growth under such challenging conditions [67]. The studies by [68] investigated the impact of various Salt-Tolerant Plant Growth-Promoting Rhizobacteria (ST-PGPR) on enhancing the productivity of salt-tolerant rice and wheat grown on sodic soils. Their findings revealed that *Lysinibacillus* sp. was particularly effective in mitigating the adverse effects of salinity. Similarly, [69] discovered the several *Bacillus* sp. as a ST-PGPR with ACC deaminase activity were the most dominant in alleviating salt stress and enhancing the biomass of rice across different agro-climatic zones. Furthermore, the siderophore-producing ability of microorganisms under stressful conditions presents a promising alternative to chemical fertilizers, potentially aiding in managing salt stress and iron limitations in salt-affected soils. [70] recently reported that salt-tolerant siderophore-producing PGPR supported rice growth and increase in protein content is associated with improved photosynthesis, which is indicative of higher chlorophyll levels.

Similarly, numerous studies have investigated the enhancement of saline tolerance in maize through various mechanisms employed by rhizobacteria to promote the growth and yield of the crop. The coinoculation of *Rhizobium* and *Pseudomonas* in *Zea mays* led to increased proline production, reduced electrolyte leakage, maintenance of leaf relative water content, and selective uptake of potassium ions, resulting in enhanced salt tolerance [47]. The [71] discovered that inoculating salt-stressed maize with *P. syringae* [72], *Enterobacter aerogenes* and *P. fluorescens* containing ACC deaminase, *Azospirillum* [73] resulted in higher K<sup>+</sup>/Na<sup>+</sup> ratios. This combination also led to elevated relative water content, chlorophyll levels, and reduced proline content [74], indicating enhanced salt tolerance mediated by various mechanisms. "The role of trehalose as an osmoprotectant

under salt stress is well-documented, with numerous ST-PGPR discovered to possess genes for trehalose biosynthetic pathways” [75-77]. Aslam et al. [78] also reported that “ACC deaminase activity in halotolerant bacterial genera such as *Arthrobacter*, *Bacillus*, *Brevibacterium*, *Gracilibacillus*, *Virgibacillus*, *Salinicoccus*, and *Pseudomonas*, as well as *Exiguobacterium* isolated from the rhizosphere and phylloplane of *Suaeda fruticosa* (L.) Forssk, stimulated the growth of maize under saline conditions”.

“The effect of ST-PGPR *S. sciuri* SAT-17 strain on anti-oxidative defense mechanisms and modulation of maize growth under salt stress was studied by” Akramet al. [79] “They reported that inoculation of maize with SAT-17 improved plant growth and decreased the ROS levels by increasing the cellular antioxidant enzyme activities (CAT, POD, and proline) under salinity treatments (75 and 150 mM NaCl). Likewise [45] studied the impact of PGPR inoculation on growth and antioxidant status of wheat under saline conditions and reported that co-inoculation with *B. subtilis* and *Arthrobacter* sp. could alleviate the adverse effects of soil salinity on wheat growth with an increase in dry biomass, total soluble sugars and proline content”. [60] studied “the effect of inoculation of *Azospirillum* strains isolated from saline or non-saline soil on yield and yield components of wheat in salinity and they observed that inoculation with the two isolates increased salinity tolerance of wheat plants; the saline-adapted isolate significantly increased shoot dry weight and grain yield under severe water salinity”. Similarly, [80] studied “the plant growth promoting activity of an auxin and siderophore producing isolate of *Streptomyces* under saline soil conditions and reported increases in growth and development of wheat plant. They observed significant increases in germination rate, percentage and uniformity, shoot length and dry weight compared to the control. Applying the bacterial inoculum increased the concentration of N, P, Fe and Mn in wheat shoots grown in normal and saline soil and thus concluded that *Streptomyces* isolate has potential to be utilized as biofertilizers in saline soils”. More recently [61] studied “the effect of five plant growth promoting halotolerant bacteria on wheat growth and found that inoculation of those halotolerant bacterial strains to ameliorate salt stress (80, 160 and 320 mM) in wheat seedlings produced an increase in root length of 71.7% in comparison with uninoculated positive controls. In particular, *Hallobacillus* sp.

and *B. halodenitrificans* showed more than 90% increase in root elongation and 17.4% increase in dry weight when compared to uninoculated wheat seedlings at 320 mM NaCl stress indicating a significant reduction of the deleterious effects of NaCl. These results indicate that halotolerant bacteria isolated from saline environments have potential to enhance plant growth under saline stress through direct or indirect mechanisms and would be most appropriate as bioinoculants under such conditions. The isolation of indigenous microorganisms from the stress affected soils and screening on the basis of their stress tolerance and PGP traits may be useful in the rapid selection of efficient strains that could be used as bioinoculants for stressed crops”.

Similarly, [45] investigated “co-inoculation with *B. subtilis* and *Arthrobacter* sp. could alleviate the adverse effects of soil salinity on wheat growth, leading to an increase in dry biomass, total soluble sugars, and proline content. [60] they observed that inoculation with these isolates of *Azospirillum* strains isolated from saline or non-saline soil increased the salinity tolerance of wheat plants, with the saline-adapted isolate significantly enhancing shoot dry weight and grain yield under severe water salinity”. The [80] explored “the plant growth-promoting activity of an auxin and siderophore-producing isolate of *Streptomyces* under saline soil conditions, including significant improvements in germination rate, shoot length, and dry weight in wheat, also increased the concentration of nitrogen, phosphorus, iron, and manganese in wheat shoots grown in normal and saline soil, indicating the potential of the *Streptomyces* isolate as a biofertilizer in saline soils”.

More recently, [61] investigated the effect of five halotolerant bacteria (*Hallobacillus* sp. and *B. halodenitrificans*) inoculation in wheat which mitigated salt stress in wheat seedlings, resulting in a significant increase in root length compared to uninoculated controls. These findings suggest that halotolerant bacteria isolated from saline environments have the potential to enhance plant growth under saline stress conditions through direct or indirect mechanisms. Utilizing indigenous microorganisms from stress-affected soils and screening them based on their stress tolerance and Plant Growth-Promoting (PGP) traits could expedite the selection of efficient strains for use as bioinoculants in stressed crops.

### 3.2 Legumes and Oil Yielding Crops

“Along with cereals, legumes maintain their significance as vital sources of protein in the human diet. Salinity poses challenges to the production of grain and food legumes in various regions worldwide. In legumes, salt stress negatively impacts root-nodule formation, symbiotic relationships, and ultimately, nitrogen fixation capacity” [81]. “The symbiotic association of rhizobia with legumes under salinity stress remains a focal area of research” [82-84]. “Many studies suggest that the application of salt-tolerant rhizobia offers a sustainable solution for enhancing the productivity of legume crops grown under salinity stress” [85]. “Several researchers have demonstrated that the adverse effects of salinity on legumes such as soybean, pigeon pea, common bean, mung bean, groundnut, and even tree legumes can be mitigated by the application of salt-tolerant rhizobial strains” [86-90]. “The role of ACC deaminase produced by Salt-Tolerant Plant Growth-Promoting Rhizobacteria (ST-PGPR) in nodule formation in legume crops is also well-documented” [91]. “During the nodulation process, ACC deaminase is crucial for enhancing the persistence of infection threads, which are negatively impacted by ethylene levels. Thus, ACC deaminase aids in nodule formation under saline conditions” [92].

“In chickpea (*Cicer arietinum* L.), delayed flowering has been directly associated with higher concentrations of Na<sup>+</sup> in the laminae of fully expanded leaves” [93]. “The inoculation of chickpea plants with *P. putida* (MSC1) or *P. pseudoalcaligens* (MSC4) isolates demonstrated an enhancement in various parameters such as phosphate solubilization, siderophore production, and IAA (indole-3-acetic acid) production, which are indicative of improved plant growth under salt stress conditions compared to uninoculated controls” [94]. Research by [95] on chickpea highlighted the significant role of Plant Growth Promoting Rhizobacteria (PGPRs) in regulating growth under salt stress. Increased concentrations of proline, malondialdehyde (MDA), as well as enhanced activities of antioxidant enzymes such as ascorbate peroxidase (APX), superoxide dismutase (SOD), and catalase (CAT) were observed under saline conditions, suggesting that inoculated PGPR strains can mitigate salinity stress by enhancing salt tolerance.

A study by [96] suggested, for the first time, the potential use of native *Pantoea dispersa* strain PSB3 as a biofertilizer to mitigate the adverse effects of salt stress on chickpea plants. *P. dispersa* exhibited notable production of IAA (218.3 µg/ml), siderophores (60.33% SU), phosphate solubilization (3.64 µg/ml), and ACC (1-aminocyclopropane-1-carboxylate) deaminase activity (207.45 nmol/mg/h) even in the presence of 150 mM NaCl under laboratory conditions. The coinoculation of ACC+ *Mesorhizobium* and rhizobacterial isolates showed more stimulatory effect on nodulation and plant biomass under normal and salt amended treatments. Results revealed that positive response of PGPR on productivity of chickpea but more enunciated response about grain yield was observed with the combined application of SA and PGPR compared to control. Growth parameters i.e root length, root mass, number of nodules and shoot mass were highly affected where SA was applied along with PGPR. From the study, it is proposed that under salt stress the combination of SA + PGPR can be a suitable practice for more production of chickpean Pakistan.

Similarly the coinoculation of 1-aminocyclopropane-1-carboxylate (ACC)-utilizing *Mesorhizobium* and rhizobacterial isolates demonstrated a more pronounced stimulatory effect on nodulation and plant biomass under both normal and salt-amended conditions [97]. The results underscored the positive impact of Plant Growth Promoting Rhizobacteria (PGPR) on chickpea productivity, with a particularly enhanced effect on grain yield observed when salicylic acid (SA) was combined with PGPR compared to control treatments. Notably, growth parameters such as root length, root mass, nodulation, and shoot mass were significantly influenced by the application of SA in conjunction with PGPR. This study suggests that the combined application of SA and PGPR could be a promising approach for enhancing chickpea production, especially in salt-stressed conditions, offering potential benefits for chickpea cultivation in [98- 101].

“The enhanced mung bean growth under saline conditions, due to bacterial inoculation, might be attributed to bacterial IAA activity, which has a tremendous effect on root growth, and water and nutrient absorption from a greater soil volume” [102]. “Inoculation/co-inoculation with rhizobia and plant growth promoting rhizobacteria (PGPR) containing 1-Aminocyclopropane-1-carboxylic acid (ACC) deaminase improve the

plant growth by reducing the stress induced ethylene production through ACC-deaminase activity” [103]. “*Cronobacter* (two isolates) and *Enterobacter* (two isolates) Inoculation of PGP bacteria under 2 and 10% salinity stress showed enhanced plant growth parameters in *Vigna radiata* compared to both salinity and non-salinity control plants” [104,105]. “Soil salinity poses a significant threat to plant health, impacting various aspects of their growth and development. Salinity disrupts the flowering and fruiting patterns of plants, leading to abnormalities in reproductive physiology, ultimately resulting in decreased crop yields and biomass. In the case of pigeon pea, salinity can cause a reduction in flowering by as much as 50% (*Cajanus cajan* L. Mill)” [106].

### 3.3 Other Crops

Salt stress profoundly impacts plant both vegetative and reproductive physiology. According to [107] in tomato plants, exposure to salinity stress leads to the accumulation of Na<sup>+</sup> in various reproductive organs such as the style, ovaries, and anther intermediate layers. This accumulation contributes to an increase in flower abortion rates, a decrease in pollen number, and a reduction in pollen viability. Additionally, high salt stress levels, such as 150 mM NaCl, can delay flowering transition and hinder the growth of shoots and roots in tomato plants.

The [108] demonstrated that certain Plant Growth Promoting Rhizobacteria (PGPR) capable of phosphate solubilization, phytohormone production, and siderophore secretion can enhance the growth of tomato plants under 2% NaCl stress conditions. Moreover, [109] found that *Bacillus velezensis* FMH2, which produces indole-3-acetic acid (IAA), significantly promotes root length and lateral root production, thereby enhancing tomato plant growth under salt stress. These findings align with the results of studies by Habib et al. [110] which demonstrated that PGPR inoculation increases the activities of reactive oxygen species (ROS)-scavenging antioxidant enzymes in okra and tomato plants under salt stress conditions.

Similarly [62] demonstrated that inoculation with *P. putida* Rs 198 promotes cotton growth and germination even under conditions of salt stress. In Arabidopsis, the impact of salinity was investigated in a hydroponic solution, revealing various symptoms such as reduced fertility,

decreased fruit length, transient wilting, and fruits predominantly containing aborted ovules and embryos, which were narrower and smaller in size [111]. Similarly, salt stress affects early flowering and the male gametophyte of canola (*Brassica napus*), resulting in a reduction in pollen grain numbers and abnormal growth of anthers, ultimately leading to decreased crop yield [112]. [113] found that under saline conditions, the growth, yield, and biomass of pearl millet are adversely affected, including reductions in germination percentage, plant height, leaf area, total biomass, and grain yield per plant. Pea plants also suffer from the adverse effects of salinity on growth, yield, and biomass [114,115] reviewed the impact of salt stress on grain legumes, noting that salinity can reduce crop yield by 12–100% in various legume species. [116] investigated the salt tolerance of black cumin (*Nigella sativa* L.) and found that increasing salinity levels from 0.3 to 9 dS m<sup>-1</sup> resulted in reduced average seed and biological yield.

Similarly, Alam et al. 2015 studied the effect of different salinity levels on the weed plant *Portulaca oleracea* L., which holds nutritional importance and is utilized similarly to spinach and lettuce in many countries. They observed reductions in biomass and yield, as well as changes in physiological attributes and alterations in stem and root structure.

## 4. SOIL REMEDIATION WITH PHYTOBENEFICIAL BACTERIA

The industrialization of the past century has led to a significant increase in the release of anthropogenic chemicals, such as Polychlorinated biphenyls (PCBs) and persistent organic pollutants (POPs), into the environment. This has resulted in detrimental effects on human health and soil ecosystems, as highlighted by Vergani et al. [117]. Soil degradation further exacerbates these issues, stemming from factors like continuous cropping, excessive use of chemical fertilizers and pesticides, and contamination by heavy metals. Microorganisms, being the predominant biota in soil, play a crucial role in restoring land ecosystems. The microecology of the rhizosphere directly or indirectly influences the growth, development, metabolic regulation, and accumulation of active ingredients in medicinal plants (MPs). [118] they suggested that, the use of microbial resources as a promising alternative to traditional fertilizers and pesticides due to their



economic efficiency, environmental safety, and non-toxic nature.

The deterioration of soil quality is accelerated by emissions from industrial waste, widespread fertilizer and pesticide use, and sewage irrigation, leading to issues like soil hardening, salinization, and accumulation of heavy metals and organic contaminants [119,120]. A national survey in China revealed that 16.1% of soil sites investigated had excessive levels of pollutants, including both inorganic and organic contaminants [121]. Despite attempts at soil amelioration through chemical and physical methods, these approaches are often inefficient, complex, and costly [122]. Therefore, there is a growing need for more effective, economical, and environmentally friendly methods and technologies to remediate degraded soils and promote sustainable ecological and agricultural development.

Rhizoremediation has emerged as a promising strategy for in situ removal of organic contaminants, facilitated by various processes performed by different species of soil bacteria. Write your own words

#### 4.1 Soil Remediation from Saline Stress through Rhizobacteria

“The presence of an excess amount of salt in soil shows cumulative and far-reaching effects on crops. Salt stress triggers ionic imbalance in plants, causes nutrient deficiency, perturbations in carbon (C) and nitrogen (N) assimilatory pathways, lowered rate of photosynthesis, generation of reactive oxygen species (ROS), osmotic and oxidative stress, thereby retarding growth and yield of crops” [123, 124].

Salt stress also poses negative impacts on soil processes, pH, decomposition rate, nutrient composition, microbial biodiversity and water availability, leading to the prevalence of drought-like conditions. According to [125] “in several agro-ecosystems, particularly in arid and semi-arid regions, drought and salinity occur simultaneously resulting in overlapping symptoms of both the stresses in the plants”. “Physical methods of treatment of saline soils, that include flushing, leaching, scraping and chemical amendments e.g. addition of gypsum and lime, are not sustainable” [126]. “These methods are time-consuming, costly and above all, cause genetic erosion of indigenous species” [127,128]. “Application of plant growth promoting rhizobacteria (PGPR) has the potential of

alleviating salt stress in plants through elicitation of several physiological and molecular mechanisms. This includes modification in root systems, inducing antioxidant machinery, production of exopolysaccharides (useful in soil aggregate formation, humification, increase in water retention, quorum sensing, nodulation and establishing microbial diversity in saline soils) and siderophores, modulation of phytohormones, synthesis of osmolytes, uptake of minerals and control of phytopathogens” [129-131]. Several species of halotolerant soil bacteria such as *Arthrobacter*, *Azospirillum*, *Alcaligenes* *Bacillus*, *Burkholderia*, *Enterobacter*, *Flavobacterium*, *Pseudomonas* and *Rhizobium*, have been reported to ameliorate salt stress in crops [132,133].

The mitigation of salt stress by halotolerant plant growth-promoting rhizobacteria (HT-PGPR) likely involves a three-tiered association: the survival of bacteria under hyperosmotic conditions, the induction of salt tolerance mechanisms in plants, and the improvement of soil quality through various mechanisms. EPS, in particular, contribute significantly to these processes by enhancing soil structure, moisture retention, and microbial interactions.

#### 4.2 HT-PGPR as Soil Ameliorators

“The involvement of microbial mechanisms in addressing saline soil through enhancements in structure and composition is equally significant. The presence of HT-PGPR in saline soil greatly influences soil quality and fertility parameters. Studies have confirmed that HTPGPR improve nutrient status, soil structure, organic matter, pH, EC, and deposition of ionic salts in soil” [134,135]. “HT-PGPR mitigate ionic toxicity through cation bridging, hydrogen bonding, and anion adsorption. There are reports where application of HT-PGPR has improved salt index of saline soil Mitigating the nutrient status, HT-PGPR improve N, C, P, Fe and Zn content of saline soils, thereby reviving the lost vegetative index and accelerating the agricultural sustainability. Under saline conditions the N content and population of nitrogen fixers are found to be decreasing. Thus, acting as an efficient reclamation strategy, the symbiotic and asymbiotic biological nitrogen fixation by salt tolerant microbes enhances the N content as well as improves fertility of soil The enrichment of saline soil using nitrogen fixing PGPR *Pseudomonas aeruginosa*, along with N compost stimulated the level of nitrogen as compared with

un-inoculated control" [136]. "Revival of arid and saline soil by utilizing salt-tolerant rhizobia can help in improving the fertility and productivity of these stressed agro-ecosystems" [137] utilized "root powder of a halophyte *Cenchrus ciliaris* as carrier to develop inoculant from HT-PGPR *B. cereus*, *P. moraviensis* and *Stenotrophomonas maltophilia*. The developed bioinoculant when applied in field improved growth of wheat and simultaneously resulted in better texture, EC, pH and organic matter of saline-sodic soil. Along with N, HT-PGPR can stimulate the P, Zn and Fe content of saline soils. *P. moraviensis* reclaimed saline sodic soil by improving P, nitrate (NO<sub>3</sub><sup>-</sup>), N and K content by almost 18– 35%" [133-137].

Microbial mechanisms play a crucial role in addressing saline soil issues by enhancing both its structure and composition. The presence of halotolerant plant growth-promoting rhizobacteria (HT-PGPR) significantly influences soil quality and fertility parameters and various aspects of soil health, including nutrient status, soil structure, organic matter content, pH levels, electrical conductivity (EC), and the deposition of ionic salts. HT-PGPR mitigate ionic toxicity through mechanisms like cation bridging, hydrogen bonding, and anion adsorption, as indicated by research conducted by Saghafi et al. [138]. Additionally, applications of HT-PGPR have been shown to enhance the salt index of saline soil, as demonstrated by.

By improving the nutrient status, HT-PGPR enhance the levels of nitrogen (N), carbon (C), phosphorus (P), iron (Fe), and zinc (Zn) in saline soils, thus revitalizing the vegetative index and promoting agricultural sustainability. In saline conditions, the nitrogen content and populations of nitrogen-fixing organisms tend to decrease. Efficient reclamation strategies involving symbiotic and asymbiotic biological nitrogen fixation by salt-tolerant microbes, as shown by Mishra et al. [139, 140] help enhance N content and improve soil fertility.

Enriching saline soil with nitrogen-fixing PGPR, such as *P. aeruginosa*, along with nitrogen compost, can stimulate nitrogen levels compared to un-inoculated controls. Innovative approaches like using root powder of halophytes as carriers for developing bioinoculants from HT-PGPR, as demonstrated by Arora et al. [136], show promise in improving soil conditions and crop growth. These bioinoculants have been shown to enhance wheat growth and improve soil texture, EC, pH, and organic matter content in saline-

sodic soil. Furthermore, HT-PGPR can stimulate the levels of phosphorus, zinc, and iron in saline soils. For instance, *P. moraviensis* has been shown to reclaim saline sodic soil by improving P, nitrate (NO<sub>3</sub><sup>-</sup>), N, and K content by significant margins, as evidenced by Hassan et al.[141].

The increase in P content of saline soil was observed by inoculation with phosphate solubilizing *B. licheniformis* MH48 strain. Reduction in soil pH, EC and enhanced availability of macro-nutrients (NPK), micronutrients (Fe, Zn, Mn and Cu) and organic matter was reported when saline soil was inoculated with HT-PGPR and phosphogypsum [142]. "Besides nutrition, aggregation is also an important soil quality which promotes water percolation, root penetration, aeration and micropore formation. The establishment of biofilm in soil aggregates or on root surface is characterized by high concentration of root exudates, signaling molecules, organic matter and water content. This complex acts as a dragging force in selecting and establishing microbial diversity. The primary content of biofilm (EPS) regulates the organic matter by serving as C source and coagulating soil particles thereby ensuring the formation of humic substances which are stable organic carbon form. Improvement of C cycling in saline soil is reported when inoculated with PGPR. Another mechanism of action reported by highlights that bacterial inoculation increases the dehydrogenase activity which is suggested to be directly correlated with soil microbial biomass which described increase in microbial biomass carbon and dehydrogenase activity in saline soil upon inoculation with HT-PGPR *B. cereus* Pb25. Research thus clearly shows the role and possible utilization of HT-PGPR in improving the quality of soils impacted with abiotic stresses such as salinity" [143-147].

HT-PGPR play a crucial role in enhancing soil structure and aggregation by producing extracellular polymeric substances (EPS), particularly under stressful conditions. This process leads to the formation of microaggregates, facilitating water percolation, root penetration, aeration, and micropore formation, as highlighted by Burns et al. [148]. The establishment of biofilms in soil aggregates or on root surfaces is characterized by a high concentration of root exudates, signaling molecules, organic matter, and water content.

This complex acts as a driving force in selecting and establishing microbial diversity.

Improving the productivity of saline soils holds the promise of not only bolstering food security but also enriching the content and quality of soil organic matter in these inherently nutrient-deficient agricultural systems. This endeavor offers multifaceted benefits in terms of soil Organic Matter Enrichment, Nutrient Retention, Food Security, Soil Structure Improvement, Carbon Sequestration and achieving the sustainability.

### 4.3 Future Prospects

A deep study on appropriate measure for Effective way to overcome traditional phytoremediation limitations when it comes to PCB removal, taking advantage also of the degrading ability of soil microorganisms and its efficiency in field application and a crucial issue to evaluate the concrete feasibility of this technology. Further detailed studies on "omic" approaches might act as powerful tools to unveil taxonomic and functional diversity of microorganisms, overcoming their limited cultivability, allowing the identification of the best plant-microbe combination involved in the saline stressed soil remediation under different environmental conditions.

An additional strategy to study salt tolerant bacteria is the exploitation of resuscitation promoting factor active on those bacteria that cannot be cultivated in vitro due to their occurrence in the soil in a viable but not culturable (VBNC) state in response to stress conditions, like high pollution levels. The resurgence promoting factor (Rpf, a cytokine from the bacterium *Micrococcus luteus*) was applied on an enrichment culture established from PCB-contaminated soil, supplying biphenyl as unique carbon source. stated that the understanding of regulatory networks of ST-PGPR in inducing salt tolerance in plants, could serve as a promising measure to alleviate salt stress and improve global food production suggested that identification of the dominant indigenous microflora from the highly saline soil and their possible adaptation mechanisms may provide a better understanding for exploring ecological and evolutionary responses in ecosystems. The role of metagenomic and metabolomic approaches becomes very important in case of harnessing and identifying novel ST-PGPR, along with the key genes and

metabolites involved in salt tolerance. especially at field scale. In particular, the potential of plants and the application of PCB-degrading endophytic bacteria remains unexplored and constitutes a big challenge due to their hydrophobic and recalcitrant chemical nature [149] which will be a in turn benefit the plant by assimilating nutrients and increasing survival/ adaption rate [150].

An in-depth exploration into overcoming the traditional limitations of phytoremediation for PCB removal involves leveraging the degrading capacity of soil microorganisms, especially in field applications. A critical aspect to assess the feasibility of this approach is to delve into "omic" techniques, which can unravel the taxonomic and functional diversity of microorganisms, circumventing their restricted cultivability. These studies, as highlighted by Canfora et al. [149], aid in identifying optimal plant-microbe combinations for remediating saline-stressed soils across various environmental conditions.

A complementary strategy involves investigating salt-tolerant bacteria, utilizing resuscitation-promoting factors to revive bacteria in a viable but non-culturable state induced by stressors like high pollution levels, as elucidated by Islam et al. [151]. Understanding the regulatory networks of salt-tolerant plant growth-promoting rhizobacteria (ST-PGPR) in enhancing plant salt tolerance, as proposed by Ramakrishnan et al. [152], offers promising avenues to mitigate salt stress and enhance global food production. Furthermore, uncovering the dominant indigenous microflora in highly saline soils and their adaptation mechanisms, as suggested by Ramakrishnan et al. [152], provides insights into ecological and evolutionary responses in ecosystems. Metagenomic and metabolomic approaches play pivotal roles in identifying novel ST-PGPR, along with the key genes and metabolites involved in salt tolerance, particularly at the field scale. However, the potential of plants and PCB-degrading endophytic bacteria remains largely untapped due to the hydrophobic and recalcitrant nature of PCBs, Nonetheless, exploring this avenue presents opportunities for plants to assimilate nutrients, thereby enhancing their survival and adaptation rates under various climatic condition [153-157].

## 5. CONCLUSION

The study aimed to assess the positive impacts of locally isolated salt-tolerant plant growth-promoting rhizobacteria (PGPR) on various crop

growth. These enhancements likely stem from the salt tolerance and growth-promoting attributes of the chosen bacterial strains. Consequently, this promising isolate could serve as a biofertilizer source, offering potential to enhance current rice cultivation methods and address salinity challenges in coastal salt-affected regions. However, it's essential to complement this initial discovery with extensive field trials for future research, ensuring suitability for large-scale implementation. Moreover, comprehensive investigations focusing on gene expression and functional traits of salt-tolerant PGPR involved in promoting plant growth under salinity stress are imperative. These studies will facilitate the development of tailored bioformulations for saline soil systems, which are increasingly prevalent worldwide. The adoption of such green biotechnology holds multifaceted positive implications for agro-ecosystems and rural environments. Revolutionizing agricultural resilience by harnessing the power of phyto-beneficial bacteria to combat saline stress and restore degraded soils, offering sustainable solutions for food security and environmental conservation.

### COMPETING INTERESTS

Authors have declared that no competing interests exist.

### COMPETING INTERESTS

Authors have declared that no competing interests exist.

### REFERENCES

1. Yadav S, Modi P, Dave A, Vijapura A, Patel D, Patel M. Effect of abiotic stress on crops. *Sustain. Crop Prod.* 2020;3(17):5-16.
2. Kumar S. Abiotic stresses and their effects on plant growth, yield and nutritional quality of agricultural produce. *Int. J. Food Sci. Agric.* 2020;4:367-378.
3. Ahmad M, Zahir ZA, Asghar HN, Asghar M. Inducing salt tolerance in mung bean through coinoculation with rhizobia and plant-growth-promoting rhizobacteria containing 1-aminocyclopropane-1-carboxylate deaminase. *Can. J. of Micro.* 2011;57(7):578-589.
4. Lambers H. Introduction: Dryland salinity: A key environmental issue in southern Australia. *Plant and Soil.* 2003;5-7.
5. Bargaz A, Lyamlouli K, Chtouki M, Zeroual Y, Dhiba D. Soil microbial resources for improving fertilizers efficiency in an integrated plant nutrient management system. *Frontiers in Microbiology.* 2018;9:1606.
6. Sultana S, Paul SC, Parveen S, Alam S, Rahman N, Jannat B, Hoque S, Rahman MT, Karim MM. Isolation and identification of salt-tolerant plant-growth-promoting rhizobacteria and their application for rice cultivation under salt stress. *Canadian Journal of Microbiology.* 2020;66(2):144-160.
7. Schirawski J, Perlin MH. Plant-microbe interaction 2017—the good, the bad and the diverse. In *International Journal of Molecular Sciences.* 2018;19(5):1374. MDPI.
8. Rütting T, Aronsson H, Delin S. Efficient use of nitrogen in agriculture. *Nutrient Cycling in Agroecosystems.* 2018;110(1):1-5.
9. Qadir M, Quilléro E, Nangia V, Murtaza G, Singh M, Thomas RJ, Drechsel P, Noble AD. Economics of salt-induced land degradation and restoration. *Natural Resources Forum.* 2014;38(4):282-295.
10. Ayyam V, Palanivel S, Chandrakasan S. *Coastal ecosystems of the tropics - adaptive management.* Springer Singapore; 2019.
11. Keren R. *Salt-affected soils, reclamation;* 2005.
12. Fita A, Rodríguez-Burruezo A, Boscaiu M, Prohens J, Vicente O. Breeding and domesticating crops adapted to drought and salinity: A new paradigm for increasing food production. *Frontiers in Plant Science.* 2015;6:978.
13. Morton MJL, Awlia M, Al-Tamimi N, Saade S, Pailles Y, Negrão S, Tester M. Salt stress under the scalpel – dissecting the genetics of salt tolerance. *The Plant Journal.* 2019;97(1):148-163.
14. Shrivastava P, Kumar R. Soil salinity: A serious environmental issue and plant growth promoting bacteria as one of the tools for its alleviation. *Saudi Journal of Biological Sciences.* 2015;22(2):123-131.
15. Patel BB, Dave RS. Studies on infiltration of saline-alkali soils of several parts of Mehsana and Patan districts of North Gujarat. *Journal of Applied Technology in Environmental Sanitation.* 2011;1(1): 87-92.

16. Bidalia A, Vikram K, Yamal G, Rao KS. Effect of salinity on soil nutrients and plant health. In M. S. Akhtar (Ed.), Salt Stress, Microbes, and Plant Interactions: Causes and Solution.. Springer Singapore. 2019;273–297.
17. Singh S, Singh V. Nutrient management in salt affected soils for sustainable crop production. Annals of Plant and Soil Research. 2022;24(2):182–193.
18. Singh K. Microbial and Enzyme Activities of Saline and Sodic Soils. Land Degradation and Development. 2016;27(3):706–718. Available:<https://doi.org/10.1002/ldr.2385>
19. Majeed A, Abbasi MK, Hameed S, Imran A, Rahim N. Isolation and characterization of plant growth-promoting rhizobacteria from wheat rhizosphere and their effect on plant growth promotion. Frontiers in Microbiology. 2015;6:198.
20. Cardinale M, Ratering S, Suarez C, Montoya AMZ, Geissler-Plaum R, Schnell S. Paradox of plant growth promotion potential of rhizobacteria and their actual promotion effect on growth of barley (*Hordeum vulgare L.*) under salt stress. Microbiological Research. 2015;181:22–32.
21. Soldan R, Mapelli F, Crotti E, Schnell S, Daffonchio D, Marasco R, Fusi M, Borin S, Cardinale M. Bacterial endophytes of mangrove propagules elicit early establishment of the natural host and promote growth of cereal crops under salt stress. Microbiological Research. 2019;223:33–43.
22. Etesami H, Beattie GA. Plant-microbe interactions in adaptation of agricultural crops to abiotic stress conditions. In Kumar V, Kumar M, Sharma S, Prasad R. (Eds.), Probiotics and Plant Health. Springer Singapore. 2017;163–200.
23. Egamberdieva D, Kucharova Z. Selection for root colonising bacteria stimulating wheat growth in saline soils. Biology and Fertility of Soils. 2009;45(6):563–571.
24. Singh RP, Jha PN. A halotolerant bacterium *Bacillus licheniformis* HSW-16 augments induced systemic tolerance to salt stress in wheat plant (*Triticum aestivum*). Frontiers in Plant Science. 2016;7:1890
25. Vurukonda SSKP, Vardharajula S, Shrivastava M, Sk ZA. Enhancement of drought stress tolerance in crops by plant growth promoting rhizobacteria. Microbiological Research. 2016;184:13–24.
26. Qurashi AW, Sabri AN. Bacterial exopolysaccharide and biofilm formation stimulate chickpea growth and soil aggregation under salt stress. Brazilian Journal of Microbiology. 2012;43:1183–1191.
27. Watanabe M, Kawahara K, Sasaki K, Noparatnaraporn N. Biosorption of cadmium ions using a photosynthetic bacterium, *Rhodobacter sphaeroides* S and a marine photosynthetic bacterium, *Rhodovulum* sp. and their biosorption kinetics. Journal of Bioscience and Bioengineering. 2003;95(4):374–378.
28. Nunkaew T, Kantachote D, Nitoda T, Kanzaki H, Ritchie RJ. Characterization of exopolymeric substances from selected *Rhodopseudomonas palustris* strains and their ability to adsorb sodium ions. Carbohydrate Polymers. 2015;115:334–341.
29. Tenney MW, Stumm W. Chemical flocculation of microorganisms in biological waste treatment. Journal Water Pollution Control Federation. 1965;1370–1388.
30. Arora M, Kaushik A, Rani N, Kaushik CP. Effect of cyanobacterial exopolysaccharides on salt stress alleviation and seed germination. Journal of Environmental Biology. 2010;31(5):701–704.
31. Chen Y, Yan F, Chai Y, Liu H, Kolter R, Losick R, Guo J. Biocontrol of tomato wilt disease by *Bacillus subtilis* isolates from natural environments depends on conserved genes mediating biofilm formation. Environmental Microbiology. 2013;15(3):848–864.
32. Ghosh D, Gupta A, Mohapatra S. A comparative analysis of exopolysaccharide and phytohormone secretions by four drought-tolerant rhizobacterial strains and their impact on osmotic-stress mitigation in *Arabidopsis thaliana*. World Journal of Microbiology and Biotechnology. 2019;35(6):90.
33. Kang SM, Joo GJ, Hamayun M, Na CI, Shin DH, Kim HY, Hong JK, Lee IJ. Gibberellin production and phosphate solubilization by newly isolated strain of *Acinetobacter calcoaceticus* and its effect on plant growth. Biotechnology Letters. 2009;31(2).
34. Richardson AE, Barea JM, McNeill AM, Prigent-Combaret C. Acquisition of

- phosphorus and nitrogen in the rhizosphere and plant growth promotion by microorganisms. *Plant and Soil*. 2009;321(1–2):305–339.
35. Yousef NM. Capability of plant growth-promoting rhizobacteria (PGPR) for producing indole acetic acid (IAA) under extreme conditions. *European Journal of Biological Research*. 2018;8(4):174–182
  36. Goswami D, Pithwa S, Dhandhukia P, Thakker JN. Delineating *Kocuria turfanaensis* 2M4 as a credible PGPR: A novel IAA-producing bacteria isolated from saline desert. *Journal of Plant Interactions*. 2014;9(1):566–576.
  37. Sandhya V, Sk ZA, Grover M, Reddy G, Venkateswarlu B. Alleviation of drought stress effects in sunflower seedlings by the exopolysaccharides producing *Pseudomonas putida* strain GAP-P45. *Biology and Fertility of Soils*. 2009;46(1):17–26.
  38. Havasi V, Hurst CO, Briles TC, Yang F, Bains DG, Hassett DJ, Sorscher E. Inhibitory effects of hypertonic saline on *P. aeruginosa* motility. In *Journal of Cystic Fibrosis*. Elsevier. 2008;7(4):267–269.
  39. Tewari S, Arora NK. Multifunctional Exopolysaccharides from *Pseudomonas aeruginosa* PF23 Involved in Plant Growth Stimulation, Biocontrol and Stress Amelioration in Sunflower Under Saline Conditions. *Current Microbiology*. 2014;69(4):484–494.
  40. Gray EJ, Smith DL. Intracellular and extracellular PGPR: commonalities and distinctions in the plant–bacterium signaling processes. *Soil Biology and Biochemistry*. 2005;37(3):395–412.
  41. Dimkpa C, Weinand T, Asch F. Plant–rhizobacteria interactions alleviate abiotic stress conditions. *Plant, Cell and Environment*. 2009;32(12):1682–1694.
  42. Lugtenberg B, Kamilova F. Plant-growth-promoting rhizobacteria. *Annual Review of Microbiology*. 2009;63(1):541–556.
  43. Dodd IC, Zinovkina NY, Safronova VI, Belimov AA. Rhizobacterial mediation of plant hormone status. *Annals of Applied Biology*. 2010;157(3):361–379.
  44. Glick BR. Bacterial ACC deaminase and the alleviation of plant stress. *Advances in Applied Microbiology*. 2004;56:291–312.
  45. Upadhyay SK, Singh JS, Saxena AK, Singh DP. Impact of PGPR inoculation on growth and antioxidant status of wheat under saline conditions. *Plant Biology*. 2012;14(4):605–611.
  46. Timmusk S, Abd El-Daim IA, Copolovici L, Tanilas T, Kännaste A, Behers L, Nevo E, Seisenbaeva G, Stenström E, Niinemets Ü. Drought-tolerance of wheat improved by rhizosphere bacteria from harsh environments: Enhanced biomass production and reduced emissions of stress volatiles. *Plos One*. 2014;9(5):e96086.
  47. Bano A, Fatima M. Salt tolerance in *Zea mays* (L). following inoculation with *Rhizobium* and *Pseudomonas*. *Biology and Fertility of Soils*. 2009;45(4):405–413.
  48. Upadhyay SK, Singh DP. Effect of salt-tolerant plant growth-promoting rhizobacteria on wheat plants and soil health in a saline environment. *Plant Biology*. 2015;17(1):288–293.
  49. Hashem A, Abd Allah EF, Alqarawi AA, Al-Huqail AA, Wirth S, Egamberdieva D. The interaction between arbuscular mycorrhizal fungi and endophytic bacteria enhances plant growth of *Acacia gerrardii* under salt stress. *Frontiers in Microbiology*. 2016;7:1089.
  50. Ali B, Wang X, Saleem MH, Sumaira, Hafeez A, Afridi MS, Khan S, Ullah I, Amaral Júnior AT do, Alatawi A. PGPR-mediated salt tolerance in maize by modulating plant physiology, antioxidant defense, compatible solutes accumulation and bio-surfactant producing genes. *Plants*. 2022;11(3):345.
  51. Shultana R, Kee Zuan AT, Yusop MR, Saud HM. Characterization of salt-tolerant plant growth-promoting rhizobacteria and the effect on growth and yield of saline-affected rice. *Plos One*. 2020;15(9):e0238537.
  52. Yan N, Marschner P, Cao W, Zuo C, Qin W. Influence of salinity and water content on soil microorganisms. *International Soil and Water Conservation Research*. 2015;3(4):316–323.
  53. Kumar H, Arora NK, Kumar V, Maheshwari DK. Isolation, characterization and selection of salt tolerant rhizobia nodulating *Acacia catechu* and *A. nilotica*. *Symbiosis*; 1999.
  54. Salwan R, Sharma A, Sharma V. Microbes mediated plant stress tolerance in saline agricultural ecosystem. *Plant and Soil*. 2019;442(1–2):1–22.

55. Misk A, Franco C. Biocontrol of chickpea root rot using endophytic actinobacteria. *Bio Control*. 2011;56(5):811–822.
56. Kruasuwan W, Thamchaipenet A. Diversity of culturable plant growth-promoting bacterial endophytes associated with sugarcane roots and their effect of growth by co-inoculation of diazotrophs and actinomycetes. *Journal of Plant Growth Regulation*. 2016;35(4):1074–1087.
57. Rungin S, Indananda C, Suttiviriya P, Kruasuwan W, Jaemsaeng R, Thamchaipenet A. Plant growth enhancing effects by a siderophore-producing endophytic streptomycete isolated from a Thai jasmine rice plant (*Oryza sativa L.* cv. KDML105). *Antonie Van Leeuwenhoek*. 2012;102(3):463–472.
58. Chen M, Wei H, Cao J, Liu R, Wang Y, Zheng C. Expression of *Bacillus subtilis* proBA genes and reduction of feedback inhibition of proline synthesis increases proline production and confers osmotolerance in transgenic *Arabidopsis*; 2007.
59. Wang W, Vinocur B, Altman A. Plant responses to drought, salinity and extreme temperatures: Towards genetic engineering for stress tolerance. *Planta*. 2003;218(1):1–14.
60. Nia SH, Zarea MJ, Rejali F, Varma A. Yield and yield components of wheat as affected by salinity and inoculation with *Azospirillum* strains from saline or non-saline soil. *Journal of the Saudi Society of Agricultural Sciences*. 2012;11(2):113–121.
61. Ramadoss D, Lakkineni VK, Bose P, Ali S, Annapurna K. Mitigation of salt stress in wheat seedlings by halotolerant bacteria isolated from saline habitats. *Springer Plus*. 2013;2(1):6.
62. Yao L, Wu Z, Zheng Y, Kaleem I, Li C. Growth promotion and protection against salt stress by *Pseudomonas putida* Rs-198 on cotton. *European Journal of Soil Biology*. 2010b;46(1):49–54.
63. Shahbaz M, Ashraf M. Improving salinity tolerance in cereals. *Critical Reviews in Plant Sciences*. 2013;32(4):237–249.
64. Roy SJ, Negrão S, Tester M. Salt resistant crop plants. *Current Opinion in Biotechnology*. 2014;26:115–124.
65. Jaemsaeng R, Jantasuriyarat C, Thamchaipenet A. Molecular interaction of 1-aminocyclopropane-1-carboxylate deaminase (ACCD)-producing endophytic *Streptomyces* sp. GMKU 336 towards salt-stress resistance of *Oryza sativa L.* cv. KDML105. *Scientific Reports*. 2018;8(1):1950.
66. Ji J, Yuan D, Jin C, Wang G, Li X, Guan C. Enhancement of growth and salt tolerance of rice seedlings (*Oryza sativa L.*) by regulating ethylene production with a novel halotolerant PGPR strain *Glutamicibacter* sp. YD01 containing ACC deaminase activity. *Acta Physiologiae Plantarum*. 2020;42(4):42.
67. Nunkaew T, Kantachote D, Kanzaki H, Nitoda T, Ritchie RJ. Effects of 5-aminolevulinic acid (ALA)-containing supernatants from selected *Rhodospseudomonas palustris* strains on rice growth under NaCl stress, with mediating effects on chlorophyll, photosynthetic electron transport and antioxidative enzymes. *Electronic Journal of Biotechnology*. 2014;17(1):19–26.
68. Damodaran T, Mishra VK, Jha SK, Pankaj U, Gupta G, Gopal R. Identification of rhizosphere bacterial diversity with promising salt tolerance, PGP traits and their exploitation for seed germination enhancement in sodic soil. *Agricultural Research*. 2019;8(1):36–43.
69. Misra S, Chauhan PS. ACC deaminase-producing rhizosphere competent *Bacillus* spp. mitigate salt stress and promote *Zea mays* growth by modulating ethylene metabolism. *3 Biotech*. 2020;10(3):119.
70. Sultana S, Alam S, Karim MM. Screening of siderophore-producing salt-tolerant rhizobacteria suitable for supporting plant growth in saline soils with iron limitation. *Journal of Agriculture and Food Research*. 2021;4:100150.
71. Nadeem SM, Zahir ZA, Naveed M, Arshad M. Preliminary investigations on inducing salt tolerance in maize through inoculation with rhizobacteria containing ACC deaminase activity. *Canadian Journal of Microbiology*. 2007;53(10):1141–1149.
72. Zerrouk IZ, Benchabane M, Khelifi L, Yokawa K, Ludwig-Müller J, Baluska F. A *Pseudomonas* strain isolated from date-palm rhizospheres improves root growth and promotes root formation in maize exposed to salt and aluminum stress. *Journal of Plant Physiology*. 2016;191:111–119.
73. Hamdia MAES, Shaddad MAK, Doaa MM. Mechanisms of salt tolerance and interactive effects of *Azospirillum brasilense* inoculation on maize cultivars

- grown under salt stress conditions. *Plant Growth Regulation*. 2004;44(2):165–174.
74. Vardharajula S, Zulfikar Ali S, Grover M, Reddy G, Bandi V. Drought-tolerant plant growth promoting *Bacillus* spp. Effect on growth, osmolytes, and antioxidant status of maize under drought stress. *Journal of Plant Interactions*. 2011;6(1):1–14.
  75. Qin S, Feng WW, Zhang YJ, Wang TT, Xiong YW, Xing K. Diversity of bacterial microbiota of coastal halophyte *limonium sinense* and amelioration of salinity stress damage by symbiotic plant growth-promoting actinobacterium glutamicibacter halophytocola KLBMP 5180. *Applied and Environmental Microbiology*. 2018;84(19):e01533-18.
  76. Orozco-Mosqueda MDC, Duan J, DiBernardo M, Zetter E, Campos-García J, Glick BR, Santoyo G. The production of ACC deaminase and trehalose by the plant growth promoting bacterium *Pseudomonas* sp. UW4 synergistically protect tomato plants against salt stress. *Frontiers in Microbiology*. 2019;10:1392.
  77. Shim JS, Seo JS, Seo JS, Kim Y, Koo Y, Do Choi Y, Jung C. Heterologous expression of bacterial trehalose biosynthetic genes enhances trehalose accumulation in potato plants without adverse growth effects. *Plant Biotechnology Reports*. 2019;13(4):409–418.
  78. Aslam F, Ali B. Halotolerant bacterial diversity associated with *Suaeda fruticosa* (L.) forssk. improved growth of maize under salinity stress. *Agronomy*. 2018;8(8):131.
  79. Akram MS, Shahid M, Tariq M, Azeem M, Javed MT, Saleem S, Riaz S. Deciphering *Staphylococcus sciuri* SAT-17 mediated anti-oxidative defense mechanisms and growth modulations in salt stressed maize (*Zea mays* L.). *Frontiers in Microbiology*. 2016;7:867
  80. Sadeghi A, Karimi E, Dahaji PA, Javid MG, Dalvand Y, Askari H. Plant growth promoting activity of an auxin and siderophore producing isolate of *Streptomyces* under saline soil conditions. *World Journal of Microbiology and Biotechnology*. 2012;28(4):1503–1509.
  81. Manchanda G, Garg N. Salinity and its effects on the functional biology of legumes. *Acta Physiologiae Plantarum*. 2008;30(5):595–618.
  82. Zahran HH. Conditions for successful *Rhizobium*-legume symbiosis in saline environments. *Biology and Fertility of Soils*. 1991;12(1):73–80.
  83. Zahran HH. *Rhizobium* -Legume Symbiosis and Nitrogen Fixation under Severe Conditions and in an Arid Climate. *Microbiology and Molecular Biology Reviews*. 1999;63(4):968–989.
  84. Graham PH. Stress tolerance in *Rhizobium* and *Bradyrhizobium*, and nodulation under adverse soil conditions. *Canadian Journal of Microbiology*. 1992;38(6):475–484.
  85. Abiala MA, Abdelrahman M, Burrett DJ, Tran LP. Salt stress tolerance mechanisms and potential applications of legumes for sustainable reclamation of salt-degraded soils. *Land Degradation and Development*. 2018;29(10):3812–3822.
  86. Bashan Y, Holguin G. *Azospirillum* – plant relationships: Environmental and physiological advances (1990–1996). *Canadian Journal of Microbiology*. 1997;43(2):103–121.
  87. Dobbelaere S, Croonenborghs A, Thys A, Ptacek D, Vanderleyden J, Dutto P, Labandera-Gonzalez C, Caballero-Mellado J, Aguirre JF, Kapulnik Y. Responses of agronomically important crops to inoculation with *Azospirillum*. *Functional Plant Biology*. 2001;28(9):871–879.
  88. Bashan Y, Holguin G, De-Bashan LE. *Azospirillum* -plant relationships: Physiological, molecular, agricultural, and environmental advances (1997-2003). *Canadian Journal of Microbiology*. 2004;50(8):521–577.
  89. Dardanelli MS, De Cordoba FJF, Espuny MR, Carvajal MAR, Díaz MES, Serrano AMG, Okon Y, Megías M. Effect of *Azospirillum brasilense* coinoculated with *Rhizobium* on *Phaseolus vulgaris* flavonoids and Nod factor production under salt stress. *Soil Biology and Biochemistry*. 2008;40(11):2713–2721.
  90. Yasin NA, Khan WU, Ahmad SR, Ali A, Ahmad A, Akram W. Imperative roles of halotolerant plant growth-promoting rhizobacteria and kinetin in improving salt tolerance and growth of black gram (*Phaseolus mungo*). *Environmental Science and Pollution Research*. 2018;25(5):4491–4505.
  91. Barnawal D, Bharti N, Maji D, Chanotiya CS, Kalra A. ACC deaminase-containing *Arthrobacter protophormiae* induces NaCl stress tolerance through reduced ACC



- oxidase activity and ethylene production resulting in improved nodulation and mycorrhization in *Pisum sativum*. *Journal of Plant Physiology*. 2014;171(11):884–894.
92. Nascimento FX, Brígido C, Glick BR, Rossi MJ. The role of rhizobial ACC deaminase in the nodulation process of leguminous plants. *International Journal of Agronomy*; 2016.
93. Pushpavalli R, Quealy J, Colmer TD, Turner NC, Siddique KHM, Rao MV, Vadez V. Salt Stress Delayed Flowering and Reduced Reproductive Success of Chickpea (*Cicer arietinum* L.), A Response Associated with Na<sup>+</sup> Accumulation in Leaves. *Journal of Agronomy and Crop Science*. 2016; 202(2).
94. Patel D, Jha CK, Tank N, Saraf M. Growth enhancement of chickpea in saline soils using plant growth-promoting rhizobacteria. *Journal of Plant Growth Regulation*. 2012;31(1):53–62.
95. Yilmaz H, Kulaz H. The effects of plant growth promoting rhizobacteria on antioxidant activity in chickpea (*Cicer arietinum* L.) under salt stress. *Legume Research-An International Journal*. 2019;42(1):72–76.
96. Panwar M, Tewari R, Gulati A, Nayyar H. Indigenous salt-tolerant rhizobacterium *Pantoea dispersa* (PSB3) reduces sodium uptake and mitigates the effects of salt stress on growth and yield of chickpea. *Acta Physiologiae Plantarum*. 2016; 38(12):278.
97. Chaudhary D, Sindhu SS. Amelioration of salt stress in chickpea (*Cicer arietinum* L.) by coinoculation of ACC deaminase-containing rhizospheric bacteria with Mesorhizobium strains. *Legume Research-An International Journal*. 2017;40(1):80–86.
98. Aneela R, Rafique M, Aftab M, Qureshi MA, Javed H, Mujeeb F, Akhtar S. Mitigation of salinity in chickpea by Plant Growth Promoting Rhizobacteria and salicylic acid. *Eurasian Journal of Soil Science*. 2019;8(3):221–228.
99. Zahir ZA, Shah MK, Naveed M, Akhter MJ. Substrate-dependent auxin production by *Rhizobium phaseoli* improves the growth and yield of *Vigna radiata* L. under salt stress conditions. *Journal of Microbiology and Biotechnology*. 2010;20(9):1288–1294.
100. Zahir ZA, Zafar-ul-Hye M, Sajjad S, Naveed M. Comparative effectiveness of *Pseudomonas* and *Serratia* sp. containing ACC-deaminase for coinoculation with *Rhizobium leguminosarum* to improve growth, nodulation, and yield of lentil. *Biology and Fertility of Soils*. 2011;47 (4):457–465.
101. Ahmad M, Zahir ZA, Khalid M, Nazli F, Arshad M. Efficacy of *Rhizobium* and *Pseudomonas* strains to improve physiology, ionic balance and quality of mung bean under salt-affected conditions on farmer's fields. *Plant Physiology and Biochemistry*. 2013;63:170–176.
102. Príncipe A, Alvarez F, Castro MG, Zachi L, Fischer SE, Mori GB, Jofré E. Biocontrol and PGPR Features in Native Strains Isolated from Saline Soils of Argentina. *Current Microbiology*. 2007;55(4):314–322.
103. Aamir M, Aslam A, Khan MY, Jamshaid MU, Ahmad M, Asghar HN, Zahir ZA. Co-inoculation with *Rhizobium* and plant growth promoting rhizobacteria (PGPR) for inducing salinity tolerance in mung bean under field condition of semi-arid climate. *Asian J Agric Biol*. 2013;1(1):7–12.
104. Desai S, Mistry J, Shah F, Chandwani S, Amaresan N, Supriya NR. Salt-tolerant bacteria enhance the growth of mung bean (*Vigna radiata* L.) and uptake of nutrients, and mobilize sodium ions under salt stress condition. *International Journal of Phytoremediation*. 2023;25(1):66–73.
105. Panwar M, Tewari R, Nayyar H. Native halo-tolerant plant growth promoting rhizobacteria *Enterococcus* and *Pantoea* sp. improve seed yield of Mungbean (*Vigna radiata* L.) under soil salinity by reducing sodium uptake and stress injury. *Physiology and Molecular Biology of Plants*. 2016P;22:445–459.
106. Garg N, Manchanda G. Effect of arbuscular mycorrhizal inoculation on salt-induced nodule senescence in *Cajanus cajan* (Pigeonpea). *Journal of Plant Growth Regulation*. 2008;27(2):115–124.
107. Ghanem ME, Van Elteren J, Albacete A, Quinet M, Martínez-Andújar C, Kinet JM, Pérez-Alfocea F, Lutts S. Impact of salinity on early reproductive physiology of tomato (*Solanum lycopersicum*) in relation to a heterogeneous distribution of toxic ions in flower organs. *Functional Plant Biology*. 2009;36(2):125–136.
108. Tank N, Saraf M. Salinity-resistant plant growth promoting rhizobacteria

- ameliorates sodium chloride stress on tomato plants. *Journal of Plant Interactions*. 2010;5(1):51–58.
109. Masmoudi F, Tounsi S, Dunlap CA, Triguí M. Endophytic halotolerant *Bacillus velezensis* FMH2 alleviates salt stress on tomato plants by improving plant growth and altering physiological and antioxidant responses. *Plant Physiology and Biochemistry*. 2021;165:217–227.
  110. Habib SH, Kausar H, Saud HM, Ismail MR, Othman R. Molecular characterization of stress tolerant plant growth promoting rhizobacteria (PGPR) for growth enhancement of rice. *Int. J. Agric. Biol*. 2016;18:184–191.
  111. Sun K, Hunt K, Hauser BA. Ovule abortion in *Arabidopsis* triggered by stress. *Plant Physiology*. 2004;135(4):2358–2367.
  112. Mahmoodzadeh H, Bemani M. Influence of salinity at early stage of flowering on the development of male gametophyte in canola (*Brassica napus L.*) cv. Symbol; 2008.
  113. Khan MA, Shaukat SS, Shahzad A, Arif H. Growth and yield responses of pearl millet (*Pennisetum glaucum [L.] R. Br.*) irrigated with treated effluent from waste stabilization ponds. *Pakistan Journal of Botany*. 2012;44(3):905–910.
  114. Wolde G, Adamu C. Impact of salinity on seed germination and biomass yields of field pea (*Pisum sativum L.*). *Asian J. Sci. Tech*. 2018;9:7565–7569.
  115. Farooq M, Gogoi N, Hussain M, Barthakur S, Paul S, Bharadwaj N, Migdadi HM, Alghamdi SS, Siddique KH. Effects, tolerance mechanisms and management of salt stress in grain legumes. *Plant Physiology and Biochemistry*. 2017;118:199–217.
  116. Faravani M, Emami DS, Gholami AB, Faravani A. The effect of salinity on germination, emergence, seed yield and biomass of black cumin. *Journal of Agricultural Sciences (Belgrade)*. 2013;58(1):41–49.
  117. Vergani L, Mapelli F, Zanardini E, Terzaghi E, Di Guardo A, Morosini C, Raspa G, Borin S. Phyto-rhizoremediation of polychlorinated biphenyl contaminated soils: An outlook on plant-microbe beneficial interactions. *Science of the Total Environment*. 2017;575:1395–1406.
  118. Wang G, Ren Y, Bai X, Su Y, Han J. Contributions of beneficial microorganisms in soil remediation and quality improvement of medicinal plants. *Plants*. 2022;11(23):3200.
  119. Sharma P, Pandey AK, Kim SH, Singh SP, Chaturvedi P, Varjani S. Critical review on microbial community during in-situ bioremediation of heavy metals from industrial wastewater. *Environmental Technology and Innovation*. 2021; 24:101826.
  120. Geng Y, Jiang L, Jiang H, Wang L, Peng Y, Wang C, Shi X, Gu J, Wang Y, Zhu J, Dai L, Xu Y, Liu X. Assessment of heavy metals, fungicide quintozene and its hazardous impurity residues in medical *Panax notoginseng* (Burk) F.H.Chen root. *Biomedical Chromatography*. 2019;33(2):e4378.
  121. Mee C, Mnr C. The report on the national soil contamination survey. Ministry of Environmental Protection; 2014.
  122. Swamy MK, Nalina N, Nalina D, Akhtar MS, Purushotham B. Heavy Metal Stress and Tolerance in Plants Mediated by Rhizospheric Microbes. In M. S. Akhtar (Ed.), *Salt Stress, Microbes, and Plant Interactions: Causes and Solution*. Springer Singapore. 2019;181–198.
  123. Bulgari R, Franzoni G, Ferrante A. Biostimulants application in horticultural crops under abiotic stress conditions. *Agronomy*. 2019;9(6):306.
  124. Mishra J, Arora NK. Secondary metabolites of fluorescent pseudomonads in biocontrol of phytopathogens for sustainable agriculture. *Applied Soil Ecology*. 2018;125:35–45.
  125. Attia H, Alamer KH, Ouhibi C, Oueslati S, Lachaâl M. Interaction between salt stress and drought stress on some physiological parameters in two pea cultivars; 2020.
  126. Egamberdieva D, Wirth S, Bellingrath-Kimura SD, Mishra J, Arora NK. Salt-tolerant plant growth promoting rhizobacteria for enhancing crop productivity of saline soils. *Frontiers in Microbiology*. 2019;10:2791.
  127. Chakraborty U, Chakraborty BN, Dey PL, Chakraborty AP. *Bacillus safensis* from wheat rhizosphere promotes growth and ameliorates salinity stress in wheat; 2018.
  128. Anderson JA, Ellsworth PC, Faria JC, Head GP, Owen MD, Pilcher CD, Shelton AM, Meissle M. Genetically engineered crops: Importance of diversified integrated pest management for agricultural sustainability. *Frontiers in Bioengineering and Biotechnology*. 2019;7:24.

129. Shahzad R, Khan AL, Bilal S, Waqas M, Kang SM, Lee IJ. Inoculation of abscisic acid-producing endophytic bacteria enhances salinity stress tolerance in *Oryza sativa*. *Environmental and Experimental Botany*. 2017;136:68–77.
130. El-Esawi MA, Alaraidh IA, Alsahli AA, Alamri SA, Ali HM, Alayafi AA. *Bacillus firmus* (SW5) augments salt tolerance in soybean (*Glycine max L.*) by modulating root system architecture, antioxidant defense systems and stress-responsive genes expression. *Plant Physiology and Biochemistry*. 2018;132:375–384
131. Arora NK, Fatima T, Mishra I, Verma M, Mishra J, Mishra V. Environmental sustainability: Challenges and viable solutions. *Environmental Sustainability*. 2018;1(4):309–340.
132. Egamberdiyeva D. Plant-growth-promoting rhizobacteria isolated from a Calcisol in a semi-arid region of Uzbekistan: Biochemical characterization and effectiveness. *Journal of Plant Nutrition and Soil Science*. 2005;168(1):94–99.
133. Yasin NA, Khan WU, Ahmad SR, Ali A, Ahmad A, Akram W. Imperative roles of halotolerant plant growth-promoting rhizobacteria and kinetin in improving salt tolerance and growth of black gram (*Phaseolus mungo*). *Environmental Science and Pollution Research*. 2018;25(5):4491–4505.
134. Vaishnav A, Shukla AK, Sharma A, Kumar R, Choudhary DK. Endophytic bacteria in plant salt stress tolerance: Current and future prospects. *Journal of Plant Growth Regulation*. 2019;38(2):650–668.
135. Meena KK, Sorty AM, Bitla UM, Choudhary K, Gupta P, Pareek A, Singh DP, Prabha R, Sahu PK, Gupta VK. Abiotic stress responses and microbe-mediated mitigation in plants: The omics strategies. *Frontiers in Plant Science*. 2017;8: 172.
136. Arora S, Vanza M. Microbial Approach for Bioremediation of Saline and Sodic Soils. In Arora S, Singh AK, Singh YP. (Eds.), *Bioremediation of Salt Affected Soils: An Indian Perspective* (pp. 87–100). Springer International Publishing; 2017.
137. Alam MA, Juraimi AS, Rafii MY, Hamid AA, Aslani F, Alam MZ. Effects of salinity and salinity-induced augmented bioactive compounds in purslane (*Portulaca oleracea L.*) for possible economical use. *Food Chemistry*. 2015;169:439–447.
138. Saghafi D, Ghorbanpour M, Shirafkan Ajirloo H, Asgari Lajayer B. Enhancement of growth and salt tolerance in *Brassica napus L.* seedlings by halotolerant *Rhizobium* strains containing ACC-deaminase activity. *Plant Physiology Reports*. 2019;24(2):225–235.
139. Mishra I, Arora NK. Rhizoremediation: A Sustainable Approach to Improve the Quality and Productivity of Polluted Soils. In NK. Arora & N. Kumar (Eds.), *Phyto and Rhizo Remediation*. Springer Singapore. 2019;9:33–66.
140. Arif MS, Shahzad SM, Yasmeen T, Riaz M, Ashraf M, Ashraf MA, Mubarak MS, Kausar R. Improving Plant Phosphorus (P) Acquisition by Phosphate-Solubilizing Bacteria. In M. Naeem, A. Ansari A, Gill SS. (Eds.), *Essential Plant Nutrients*. Springer International Publishing. 2017; 513–556.
141. Hassan TU, Bano A, Naz I. Halophyte root powder: An alternative biofertilizer and carrier for saline land. *Soil Science and Plant Nutrition*, 2018;64(5):653–661.
142. Sandhya V, Ali Sk Z. The production of exopolysaccharide by *Pseudomonas putida* GAP-P45 under various abiotic stress conditions and its role in soil aggregation. *Microbiology*. 2015;84 (4):512–519.
143. Arora S, Singh YP, Vanza M, Sahni D. Bioremediation of saline and sodic soils through halophilic bacteria to enhance agricultural production. *Journal of Soil and Water Conservation*. 2016;15(4): 302–305.
144. Rashid MI, Mujawar LH, Shahzad T, Almeelbi T, Ismail IM, Oves M. Bacteria and fungi can contribute to nutrients bioavailability and aggregate formation in degraded soils. *Microbiological Research*. 2016;183:26–41.
145. Verma M, Mishra J, Arora NK. Plant growth-promoting rhizobacteria: Diversity and applications. In Sobti RC, Arora NK, Kothari R. (Eds.), *Environmental Biotechnology: For Sustainable Future*. Springer Singapore. 2019;129–173.
146. Ul Hassan T, Bano A. Construction of IAA-Deficient Mutants of *Pseudomonas moraviensis* and Their Comparative Effects with Wild Type Strains as Bio-inoculant on Wheat in Saline Sodic Soil. *Geomicrobiology Journal*. 2019;36(4):376–384.

147. Al-Enazy AA, Al-Barakah F, Al-Oud S, Usman A. Effect of phosphogypsum application and bacteria co-inoculation on biochemical properties and nutrient availability to maize plants in a saline soil. *Archives of Agronomy and Soil Science*. 2018;64(10):1394–1406.
148. Burns RG, DeForest JL, Marxsen J, Sinsabaugh RL, Stromberger ME, Wallenstein MD, Weintraub MN, Zoppini A. Soil enzymes in a changing environment: Current knowledge and future directions. *Soil Biology and Biochemistry*. 2013;58:216–234.
149. Canfora L, Bacci G, Pinzari F, Lo Papa G, Dazzi C, Benedetti A. Salinity and bacterial diversity: To what extent does the concentration of salt affect the bacterial community in a saline soil? *Plos One*. 2014;9(9):e106662..
150. Lipińska A, Wyszowska J, Kucharski J. Diversity of organotrophic bacteria, activity of dehydrogenases and urease as well as seed germination and root growth *Lepidium sativum*, *Sorghum saccharatum* and *Sinapis alba* under the influence of polycyclic aromatic hydrocarbons. *Environmental Science and Pollution Research*. 2015;22(23):18519–18530.
151. Islam S, Akanda AM, Prova A, Islam MT, Hossain MM. Isolation and identification of plant growth promoting rhizobacteria from cucumber rhizosphere and their effect on plant growth promotion and disease suppression. *Frontiers in Microbiology*. 2016;6:1360.
152. Ramakrishnan B, Maddela NR, Venkateswarlu K, Megharaj M. Potential of microalgae and cyanobacteria to improve soil health and agricultural productivity: A critical view. *Environmental Science: Advances*. 2023;2(4):586–611.
153. Rillig MC, Muller LA, Lehmann A. Soil aggregates as massively concurrent evolutionary incubators. *The ISME Journal*. 2017;11(9):1943–1948.
154. Vaishnav A, Shukla AK, Sharma A, Kumar R, Choudhary DK. Endophytic bacteria in plant salt stress tolerance: Current and future prospects. *Journal of Plant Growth Regulation*. 2019;38:650-668.
155. Khan MA, Asaf S, Khan AL, Ullah I, Ali S, Kang SM, Lee IJ. Alleviation of salt stress response in soybean plants with the endophytic bacterial isolate *Curtobacterium* sp. SAK1. *Annals of Microbiology*. 2019;69(8):797–808.
156. Ma Y, Prasad MNV, Rajkumar M, Freitas H. Plant growth promoting rhizobacteria and endophytes accelerate phytoremediation of metalliferous soils. *Biotechnology Advances*. 2011;29(2):248–258.
157. Kim K, Samaddar S, Chatterjee P, Krishnamoorthy R, Jeon S, Sa T. Structural and functional responses of microbial community with respect to salinity levels in a coastal reclamation land. *Applied Soil Ecology*. 2019;137:96–105.

© Copyright (2024): Author(s). The licensee is the journal publisher. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:  
<https://www.sdiarticle5.com/review-history/114466>