



# Effect of Supplementing Wheat Bran, Cowpea (*Vigna unguiculata*) Hay, and Their Combinations to Begait Lambs Fed Natural Grass Hay as a Base Diet on Their Nutrient Digestibility and Dry Matter

Gebreslasie Gebrekidan Beyene <sup>a\*</sup>

<sup>a</sup> Humera Begait Animal Research Center, P. O. Box 62, Humera, Ethiopia.

## Author's contribution

The sole author designed, analysed, interpreted and prepared the manuscript.

## Article Information

### Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/123661>

Original Research Article

Received: 18/07/2024

Accepted: 20/09/2024

Published: 02/10/2024

## ABSTRACT

### Aim/Objective:

- To evaluate the effect of supplementing wheat bran (WB), cowpea hay (CPH), and mixes of the two on dry matter and nutrient digestibility of Begait lambs fed natural grass hay as a basal diet.
- To determine the ideal ratio for supplementing wheat bran and cowpea hay on the dry matter and nutritional digestibility of Begait lambs given natural grass hay as their baseline diet.

\*Corresponding author: E-mail: [gebre077@gmail.com](mailto:gebre077@gmail.com);

**Cite as:** Beyene, Gebreslasie Gebrekidan. 2024. "Effect of Supplementing Wheat Bran, Cowpea (*Vigna Unguiculata*) Hay, and Their Combinations to Begait Lambs Fed Natural Grass Hay As a Base Diet on Their Nutrient Digestibility and Dry Matter". *Asian Journal of Research in Animal and Veterinary Sciences* 7 (4):343-52. <https://www.journalajravs.com/index.php/AJRAVS/article/view/318>.

**Study Design:** The study employed a randomized complete block design (RCBD) with five treatments and blocks.

**Location and Length of Study:** Northern Ethiopia, Humera Agricultural Research Center farm, starting December 2016 until November 2017.

**Methodology:** 25 intact male Begait lambs were bought from the local market, weighing an average of  $28.02 \pm 1.49$  kg (mean  $\pm$  SD) at 5 to 6 months of age. Based on their starting body weight, they were split into five groups and assigned randomly to each of the five treatments. The standard diet for T1, T2, T3, T4, and T5 consisted of giving natural grass hay *ad libitum* plus supplements of 300 g CPH, 225 g CPH + 75 g WB, 150 g CPH + 150 g WB, 75 g CPH + 225 g WB, and 300 g WB. DM day-1. All lambs in each treatment were equipped with a fecal-collection bag for three days, after which they were required to collect all of their excrement for seven days to measure nutrient use.

**Results:** Among the treatment groups, differences in DM, CP ( $P < .001$ ), OM, and ADF ( $P < .01$ ) digestibility were noted. T5 and T3 showed higher ( $P < .001$ ) DM and CP digestibility, respectively, in comparison to the other treatment groups. There was no difference in the digestibility of ADF between T5 and T3, but it was higher for T5 than for the lambs in T1, T2, and T4. Conversely, T1 showed low digestion of OM and CP.

**Conclusion:** Thus, the study concluded that supplementing Begait lambs with an equal amount of cowpea to wheat bran (150 g CPH + 150 g WB) outweighs the dry matter and nutritional digestibility of the lambs and that it is crucial to address the feed scarcity in the study area at a reasonable cost.

**Keywords:** Begait lamb; digestibility; *Vigna unguiculata*; wheat bran.

## 1. INTRODUCTION

In the next few years, the global population will be expected to rise. The demand for cattle and livestock products will rise as a result. To address this demand, effective and sustainable feeding systems must be developed.

The main source of feed for ruminants in tropical regions is rangelands with diverse native vegetation, which includes grasses, legumes, trees, and shrubs. Ruminants rely on perennial grasses and agricultural byproducts, such as crop leftovers and agro-industrial byproducts, when pasture supply declines during dry seasons. Crop leftovers are fibrous remnants whose low digestion, low voluntary intake, and low levels of nitrogen, calories, minerals, and vitamins restrict their feeding usefulness [1]. Similar to this, forage quality and availability are unfavorable throughout the year, meaning that gains achieved during the wet season are either completely or partially lost during the dry season [2]. Thus, attempts have been made to increase the nutritional content of these fibrous feeds. Herbaceous legume forages could be one way to supplement.

The Humera Agricultural Research Center (HuARC) introduced several improved herbaceous legume forages, such as cowpea, lablab, and clitoria, to the area as intervention measures. Temesgen, a cowpea variety, was listed in the national variety release in 2014 G.C.

as the best feed resource in the region and among other comparable agroecologies in the nation, based on monitoring and evaluation of the varieties [3]. It is a valuable crop because it serves two purposes: human food and animal fodder.

Temesgen is distinguished by its high crude protein content (17.15% CP) and high herbage DM output ( $11.9 \text{ t ha}^{-1}$ ). Because of its high protein content, it may be used to replace the protein that ruminants consume on roughages of lower quality [4, 5], including matured natural pasture hay [6]. Animal feeding trials were necessary to ensure appropriate consumption because the importance of improving the digestibility of nutrients and dry matter is not given much attention in the existing literature. Thus, the following goals guided the conduct of this study: the objectives are:

- To ascertain the impact of supplementing wheat bran, cowpea (*Vigna unguiculata*) hay, and their combinations on the dry matter and nutritional digestibility of Begait lambs fed natural grass hay as their basal diet;
- To determine the ideal ratio for supplementing wheat bran and cowpea hay on the dry matter and nutritional digestibility of Begait lambs, provide natural grass hay as their fundamental diet.

## 2. MATERIALS AND METHODS

### 2.1 Description of the Study Area

The Humera Agricultural Research Center (HuARC), located in the western zone of Tigray, Northern Ethiopia, served as the study's on-station location. It is located at latitude 14°15' N and longitude 36°37' E, at an altitude of 608.9 m.a.s.l. Vertisol is the most common soil type. The climate at the location is semi-arid with a unimodal rainfall pattern. Mean maximum and minimum temperatures ranged from 33 to 41.7 and 17.5 to 22.2°C, respectively, and the mean annual rainfall was 581.2 mm, which was received during the summer/rainy season [7].

### 2.2 Experimental Animals' Management

Twenty-five intact male Begait lambs, aged between five and six months and weighing an average of 28.02 ± 1.49 kg (mean ± SD), were purchased at the nearby market. The owners gave the age information, which was confirmed through dentition. They received vaccinations against common diseases, internal parasites, and external parasites upon arriving at the center. They were also dewormed and sprayed. The 90-day stay of the lambs was preceded by a 15-day period during which they were acclimated to their new surroundings and food while considering animal welfare. During the experimental time, the lambs were housed in separate pens with plastic buckets for hydration and supplements and feeding troughs for hay.

### 2.3 Experimental Design and Treatments

Five blocks of each of the five treatments were employed in a randomized complete block design (RCBD) (Table 1). At the start of the feeding trial, lambs were grouped based on their initial body weight, which was calculated as the average of two subsequent weighings following an overnight fast. Each block's lambs were randomized to receive one of the five treatment meals. They were placed in separate pen that measured 115 cm in length and 85 cm in width in a shed that had natural ventilation. Every lamb had unrestricted access to fresh, clean water and common salt. The ten-day trial was divided into three days for the suitability of the fecal-collecting bag and seven days for the actual feces collection.

The treatment setup was based on the suggestions of earlier authors [8] and [9], who

supplemented 300 g of cowpea haulms and Desmodium hay to achieve substantial effects on growth performance, DM intake, and digestibility on sheep, respectively.

### 2.4 Digestibility Trial

To evaluate the use of various dietary components, all lambs engaged in a digestibility trial at the end of the feeding trial. This experiment was conducted using the same lambs and rations as the feeding trial. Every lamb in each treatment was equipped with a fecal collection bag for three days, after which they had to collect all of their excrement for seven days. Every day, specimens and records of food given and rejected were compiled. Every animal's daily fecal excretion was gathered and noted. Each animal's daily fecal output was measured by emptying the bag before providing food and drink in the morning. Each animal's excrement was carefully mixed, with 20% of it being sampled every day and kept in labeled deep freezers at -20°C.

Throughout the collecting period, the samples were put together for each animal, and 20% of the composite sample was taken out, weighed, and partially dried at 60°C for 72 hours in a forced draft oven to achieve a constant weight. After being ground to fit through a 1 mm filter, the dry fecal samples were sealed in polyethylene bags and set aside for additional chemical examination. Ultimately, each lamb's apparent digestibility coefficients (ADC) were determined using the guidelines provided in [10]:

$$\text{ADC (\%)} = \frac{((\text{nutrient intake} - \text{fecal excretions}) / \text{nutrient intake}) \times 100}{(1)}$$

### 2.5 Chemical Analysis

To ascertain nutritional content, representative samples of feed and excrement were taken to the animal nutrition laboratory center at Haramaya University. As per the guidelines provided by the Association of Official Analytical Chemists [11], analysis was conducted for DM, ash, and N concentrations. By igniting in a muffle furnace at 550 degrees Celsius for six hours and oven drying at 105 degrees Celsius for a whole night, respectively, the ash and dry matter contents were ascertained.

Crude protein (CP) was calculated as  $N \times 6.25$ , and the nitrogen (N) concentration was

**Table 1. Experimental Treatments**

Treatment	Grass hay	Amount of CPH supplement (g)	Amount of WB supplement (g)	Supplement amount (g head <sup>-1</sup> day <sup>-1</sup> )	CP (g)	ME (MJ)
1	Ad libitum	300	0	300	51.4	6.7
2	Ad libitum	225	75	300	50.9	7.9
3	Ad libitum	150	150	300	50.5	7.9
4	Ad libitum	75	225	300	49.9	7.2
5	Ad libitum	0	300	300	49.5	8.2

CPH: cow pea hay; WB: wheat bran; CP: crude protein; ME: metabolizable energy

**Table 2. Chemical composition of experimental feeds**

Feed offered	DM %	CP	NDF	ADF	ADL	Ash
				DM %		
Cowpea hay	88.75	17.15	60.00	50.30	14.50	12.50
WB	88.22	16.50	38.46	16.35	5.75	6.45
Grass hay	90.65	5.46	77.15	53.28	14.76	9.13

CP: crude protein; NDF: neutral detergent fiber; ADF: acid detergent fiber; ADL: acid detergent lignin; DM: dry matter; WB: wheat bran

ascertained using the Kjeldahl technique. The methods of [12] were used to determine the amounts of acid detergent fiber (ADF), neutral detergent fiber (NDF), and acid detergent lignin (ADL). The formula for organic matter was 100 minus ash.

## 2.6 Partial Budget Analysis

Using the method of [13], a partial budget analysis was carried out to assess the financial benefit of the different treatment options. It had a role in the benefits and variable costs computation. Total return (TR) in the analysis was defined as the difference in lamb selling prices for each treatment before and after the trial. For the computation of the variable expenses, the expenditures spent on various feedstuffs were taken into consideration. The actual feed intake for the duration of the feeding period was multiplied by the current price to determine the cost of the feeds.

The partial budget technique computes net return (NR), or the amount of money remaining after deducting total variable costs (TVC) from total returns (TR), and measures profit or losses, which are the differences between gains and expenses for the proposed change:

$$NR = TR - TVC \quad (2)$$

The total cost of variable costs is the sum of all input costs that vary as a result of technology

used for production changes. As a reference criterion for deciding whether to adopt new technology, the change in net return ( $\Delta NR$ ) was computed by subtracting the change in total variable cost ( $\Delta TVC$ ) from the change in total return ( $\Delta TR$ ).

$$\Delta NR = \Delta TR - \Delta TVC \quad (3)$$

The rise in net income ( $\Delta NR$ ) corresponding to each increased unit of expenditure ( $\Delta TVC$ ) is measured by the marginal rate of return (MRR). A percentage is used to express this.

$$MRR\% = (\Delta NR / \Delta TVC) \times 100 \quad (4)$$

## 2.7 Analytical Statistics

Using the Generalized Linear Model (GLM) of the Statistical Analysis System for Windows (SAS, 2002), data on apparent digestibility coefficients were subjected to the analyses of variance (ANOVA) technique by a randomized complete-block design to detect treatment effects. A Fisher's Least Significant Difference (LSD) test was used to compare treatment means at a 5% probability level. The experiment was analyzed using the following statistical model:

$$Y_{ij} = \mu + T_i + B_j + e_{ij} \quad (5)$$

Where:  $Y_{ij}$  = response variable;  $\mu$  = the overall mean;  $T_i$  = the  $i$ th treatment effect;  $B_j$  = the  $j$ th block effect (initial BW);  $e_{ij}$  = the random error

**Table 3. Apparent digestibility coefficients of dry matter and nutrients of Begait lambs fed natural grass hay and supplemented with *Vigna unguiculata* hay, wheat bran, and their mixes**

Digestibility (%)	Treatments					SL	PSE
	T1	T2	T3	T4	T5		
DM	61.36 <sup>b</sup>	61.29 <sup>b</sup>	62.69 <sup>b</sup>	61.57 <sup>b</sup>	69.94 <sup>a</sup>	***	0.981
OM	60.75 <sup>b</sup>	68.45 <sup>a</sup>	67.73 <sup>a</sup>	69.05 <sup>a</sup>	70.15 <sup>a</sup>	**	0.988
CP	58.84 <sup>c</sup>	66.06 <sup>b</sup>	70.87 <sup>a</sup>	62.92 <sup>b</sup>	65.15 <sup>b</sup>	***	1.139
NDF	60.10	65.53	65.08	66.21	66.41	ns	2.554
ADF	53.89 <sup>b</sup>	52.65 <sup>b</sup>	54.16 <sup>ab</sup>	50.68 <sup>c</sup>	55.79 <sup>a</sup>	**	0.511

*a-c, means with different superscripts in a row are significantly different; \*\*\* = (p<.001); \*\* = (p<.01); ns: non-significant; PSE: pooled standard error of the mean; SL: level of significance; CP: crude protein; NDF: neutral detergent fiber; ADF: acid detergent fiber; DM: dry matter; OM: organic matter; T1: received 300g cowpea hay; T2: received 225g cowpea hay + 75g wheat bran; T3: received 150g cowpea hay + 150g wheat bran; T4: received 75g cowpea hay + 225g wheat bran; T5: received 300g wheat bran*

### 3. RESULTS AND DISCUSSION

#### 3.1 Digestibility of Nutrients and Dry Matter

Table 3 shows that there were differences in the digestibility of DM, CP ( $P<.001$ ), OM, and ADF ( $P<.01$ ) across the treatment groups. T1 revealed low digestion of OM and CP. T5 and T3 showed higher ( $P<.001$ ) DM and CP digestibility, respectively, in comparison to the other treatment groups. There was no difference in the digestibility of ADF between T5 and T3; however, it was higher for T5 than for the lambs in T1, T2, and T4.

The lowered NDF and ADF contents of WB (Table 2) and increased CP consumption (Table 4) may be the cause of the higher DM and CP digestibility seen in T5 and T3, respectively. The combination of the two supplements in similar amounts may have promoted high rumen fermentation, permitted good synthesis of volatile fatty acids (VFAs) for absorption, and boosted rumen biomass production, as suggested by the higher CP digestibility in T3 [14]. Conversely, a lower CP intake (Table 4) may be the cause of the low digestion of DM and ADF seen in T4.

The obtained DM digestibility values are between 60 and 69%, which is considered to be an indication of a reasonably high digestible level [15]. Similar levels of nutrient availability from the various combinations of the treatment diets were indicated by the lack of a significant variation in NDF digestibility in the current investigation. Previous reports of comparable outcomes exist [16–18]. Furthermore, [19] observed that there was no noticeable distinction in the digestibility of NDF and ADF, which was explained by a

potential decrease in rumen pH caused by giving the animals more digestible supplements, which in turn hurts the number of rumen bacteria that ferment cell walls. Microorganism activity in the rumen is necessary for digestion. For the bacteria to function properly, they also need energy, nitrogen, minerals, and an appropriate medium [20].

#### 3.2 Analysis of a Partial Budget

The lambs fed with 150 g cowpea hay and 150 g wheat bran (T3) exhibited a better net return and a change in net return, followed by 225 g cowpea hay and 75 g wheat bran (T2) (Table 5). The net return of lambs fed 225 g WB and 75 g cowpea hay (T4) was poor. The improved growth and physical state of the lambs may be the cause of the higher net return and net return change seen in T3. Conversely, a higher total variable cost may be the cause of the lower net return seen in T4.

The cost of supplementing with wheat bran was higher than that of a diet based on cowpea hay. The primary expenses associated with the cowpea hay-based treatment were clearing the ground, planting, weeding, harvesting, and chopping. Comparatively, the partial budget analysis showed that T3 had a greater MRR than the others, which would have resulted from the lambs' higher selling price. The marginal rate of return evaluates the increase in net income and the effects of extra investment in new technology on additional net return. Therefore, data showed that giving wheat bran as a supplement caused an economic loss, whereas providing an extra unit of one birr per lamb cost increment resulted in a 0.50- and 1.25-birr advantage for T2 and T3, respectively.

**Table 4. Daily dry matter and nutrient intake of Begait lambs fed natural grass hay and supplemented with *Vigna unguiculata* hay, wheat bran, and their mixes**

Intake (gday <sup>-1</sup> )	T1	T2	T3	T4	T5	SL	PSE
Grass hay DM	387.11 <sup>b</sup>	369.41 <sup>b</sup>	419.41 <sup>a</sup>	330.99 <sup>c</sup>	373.47 <sup>b</sup>	***	6.801
Supplement DM	300.00	300.00	300.00	300.00	300.00	ns	0.000
Total DM	687.11 <sup>b</sup>	669.41 <sup>b</sup>	719.41 <sup>a</sup>	630.99 <sup>c</sup>	673.47 <sup>b</sup>	***	6.801
Total DM (%BW)	2.21 <sup>a</sup>	2.08 <sup>by</sup>	2.17 <sup>ab</sup>	2.01 <sup>c</sup>	2.11 <sup>b</sup>	**	0.099
Nutrient							
Total OM	682.07 <sup>a</sup>	669.35 <sup>bc</sup>	723.27 <sup>a</sup>	641.55 <sup>c</sup>	689.30 <sup>ab</sup>	**	7.625
Total CP	84.800 <sup>ab</sup>	82.631 <sup>ab</sup>	85.38 <sup>a</sup>	79.218 <sup>c</sup>	82.338 <sup>b</sup>	**	0.599
Total NDF	526.79 <sup>a</sup>	491.04 <sup>a</sup>	514.64 <sup>a</sup>	422.66 <sup>b</sup>	445.37 <sup>b</sup>	***	9.754
Total ADF	400.50 <sup>a</sup>	358.35 <sup>b</sup>	353.70 <sup>b</sup>	277.08 <sup>c</sup>	279.21 <sup>c</sup>	***	11.109
ME(MJday <sup>-1</sup> )	6.74 <sup>c</sup>	7.91 <sup>ab</sup>	7.85 <sup>ab</sup>	7.24 <sup>bc</sup>	8.17 <sup>a</sup>	**	0.160

a-c, means with different superscripts in a row are significantly different; \*\*\*= (p<.001); \*\*= (p<.01); PSE: pooled standard error of mean; SL: level of significance; ME: metabolizable energy; ns: not significant difference; CP: crude protein; NDF: neutral detergent fiber; ADF: acid detergent fiber; DM: dry matter; OM: organic matter; T1: received 300g cowpea hay; T2: received 225g cowpea hay + 75g wheat bran; T3: received 150g cowpea hay + 150g wheat bran; T4: received 75g cowpea hay + 225g wheat bran; T5: received 300g wheat bran.

Source: [7]

**Table 5. Partial budget analysis of Begait lambs fed natural grass hay and supplemented with *Vigna unguiculata* hay, wheat bran, and their mixtures**

Variables (ETB)	Treatments				
	T1	T2	T3	T4	T5
Animal count	5	5	5	5	5
Lambs purchase price (ETB head <sup>-1</sup> )	800	800	800	800	800
Total intake of the basal diet (kg head <sup>-1</sup> )	37.58	35.90	40.41	31.91	36.15
Total intake of cowpea hay (kg head <sup>-1</sup> )	30.42	22.82	15.21	7.61	0
Total intake of wheat bran (kg head <sup>-1</sup> )	0	7.65	15.3	22.95	30.6
Basal diet expenditure (ETB head <sup>-1</sup> )	27.43	26.21	29.49	23.29	26.39
Cost of cowpea (ETB head <sup>-1</sup> )	75.75	56.82	37.87	18.95	0
Cost of wheat bran (ETB head <sup>-1</sup> )	0	26.77	53.55	80.32	107.1
Total variable cost (ETB head <sup>-1</sup> )	103.18	109.81	120.92	122.57	133.49
Δ TVC	0	6.63	17.74	19.39	30.31
Selling price of sheep (ETB head <sup>-1</sup> )	1060	1070	1100	1040	1090
Total Rate of Return (TRR)	260	270	300	240	290
ΔTRR	0	10	40	-20	30
Net return (NR) ETB head <sup>-1</sup> )	156.82	160.19	179.08	117.43	156.51
ΔNR	0	3.37	22.26	-39.39	-0.312
Marginal rate of return (MRR, %)	0	50.83	125.48	-203.15	-1.03

ETB: Ethiopian Birr; TVC: total variable cost; T1: 300g cowpea hay; T2: 225g cowpea hay + 75g wheat bran; T3: 150g cowpea hay + 150g wheat bran; T4: 75g cowpea hay + 225g wheat bran; T5: 300g wheat bran

#### 4. CONCLUSION

This study revealed that food had an impact on the dry matter and nutrient digestibility of Begait lambs except NDF. Lambs fed 300 g of wheat bran and 150 g CPH + 150 g WB, respectively, exhibited higher levels of DM and CP digestibility. Thus, the study concluded that supplementing Begait lambs with an equal amount of cowpea to wheat bran (150 g CPH + 150 g WB) outweighs the dry matter and nutritional digestibility of the lambs and that it is crucial to address the feed scarcity in the study area at a reasonable cost.

#### DISCLAIMER (ARTIFICIAL INTELLIGENCE)

The author hereby declares that NO generative AI technologies such as large language models (ChatGPT, COPILOT, etc.) and text-to-image generators have been used during the writing or editing of this manuscript.

#### ACKNOWLEDGEMENTS

For financial support, the author is grateful to the Humera Agricultural Research Center (HuARC) and the Tigray Agricultural Research Institute (TARI). In addition, the author thanks Haramaya

University Nutrition Laboratory Center for contributing the services necessary to perform the feed and excrement chemical analyses.

### ETHICAL APPROVAL

"The author hereby declares that "Principles of Laboratory Animal Care" (NIH publication No. 85-23, revised 1985) were followed, as well as specific national laws where applicable. All experiments have been examined and approved by the appropriate ethics committee."

### COMPETING INTERESTS

The author has declared that no competing interests exist.

### REFERENCES

1. Mengistu A, Kebede G, Feyissa F, Assefa G. Review on major feed resources in Ethiopia: conditions, challenges, and opportunities. *Academic Research Journal of Agricultural Science and Research*. 2017;5(3):176-185.
2. Alemayehu M. Country pastures/forage resource profiles. Ethiopia. Food and Agriculture Organization of the United Nations (FAO); 2006.
3. Solomon G, Kibrom G. Forage production potential and nutritive value of cowpea (*Vigna unguiculata*) genotypes in the northern lowlands of Ethiopia. *E3 Journal of Agricultural Research and Development*. 2014; 5(4): 066-071.
4. Ravhuhali KE, JW Ng'ambi, D Norris, VI Ayodele. The feeding value of four cowpea hay cultivars and the effect of their supplementation on intake and digestibility of buffalo grass hay fed to Pedi goats. 2011; 909-922.
5. Gwanzura T, JW Ng'ambi, D Norris. Nutrient composition and tannin contents of forage sorghum, cowpea, lablab, and Mucuna hays grown in Limpopo province of South Africa. 2012; 256-262.
6. Baloyi JJ, Ngongoni NT, Hamudikuwanda H. Chemical composition and ruminal degradability of cowpea and silver leaf *Desmodium* forage legumes harvested at different stages of maturity. *Tropical and Subtropical Agro-ecosystems*. 2008;8:81-91
7. Gebrekidan G, Teklebrhan T, Tesfay Z. Growth performance and carcass traits of Begait lambs fed diets of cowpea (*Vigna unguiculata*) hay, wheat bran, and their mixtures. *Journal of Agriculture and Ecology Research International*. 2019;20(1):1-12.
8. Korlagama KDN, Mould FL, Fernandez-Rivera S, Hanson J. The effect of supplementing maize Stover with cowpea (*Vigna unguiculata*) haulms on the intake and growth performance of Ethiopian sheep. *Animal*. 2008; 2(6): 954–961.
9. Tolera A, Sundstøl F. Supplementation of graded levels of *Desmodium intortum* hay to sheep feeding on maize stover harvested at three stages of maturity: 1. Feed intake, digestibility, and body weight change. *Animal Feed Science and Technology*. 2000; 85(3): 239-257.
10. Asad F, Rehman T, Qureshi NA, Tahir N. Estimation of apparent digestibility coefficient of plant feed ingredients (soybean and sunflower meal) for Labeo Rohita. *American Journal of Biomedical and Life Sciences*. 2013;1(1):8-11.
11. Association of Official Analytical Chemists (AOAC). Official methods of analysis, 18th edition. Association of Official Analytical Chemists, Maryland, USA; 2005.
12. Van Soest PJ, Robertson BJ. Analysis of forage and fibrous feeds. A laboratory manual for animal science (613). Cornell University, Ithaca, New York, USA; 1985.
13. Upton M. Farm management in Africa: the principle of production and planning. Oxford University Press, Great Britain. 1979;282
14. McDonald P, Edwards RA, Greenhalgh JFD, Morgan CA. *Animal Nutrition*, 5<sup>th</sup> edition. Longman Publishers Ltd., Singapore. 1995;583-585.
15. Rinehart L. Ruminant nutrition for grazers. ATTRA–National Sustainable Agriculture Information Service;2008.
16. Alemu W, Melaku S, Tolera A. Supplementation of cottonseed, linseed, and noug seed cakes on feed intake, digestibility, body weight, and carcass parameters of Sidama goats. *Tropical animal health and production*. 2010; 42(4): 623-631.
17. Dessie Jemberu, Solomon Melaku, Firew Tegegne, Kurt J Peters. Effect of supplementation of Simada sheep with graded levels of concentrate meal on feed intake, digestibility, and body-weight parameters. *Tropical animal health and production*. 2010; 42: 841-848.

18. Birhanu Tesema, Getachew Animut, Mengistu Urge. Effect of green *prosopis juliflora* pods and noug seed (*Guizotia obissynica*) cake supplementation on digestibility and performance of Blackhead Ogaden sheep fed hay as a basal diet. Science, Technology, and Art Research Journal. 2013; 2(2): 38-47.
19. Solomon M, Melaku S, Tolera A. Supplementation of cottonseed meal on feed intake, digestibility, and live weight and carcass parameters of Sidama goats. Livestock Science. 2008; 119 (1): 137-144.
20. Ranjhan SK. Animal nutrition in the tropics, 5th Edition. Vikas Publishing House PVT Ltd., New Delhi, India. 2001; 556-559.



## APPENDIX

### List of Tables in the Appendix:

**Appendix Table 1. Summary of ANOVA for apparent digestibility of dry matter and nutrients of experimental lambs**

Digestibility	DF	MS	F-value	Pr >F	SL	CV (%)
DM	10.0	41.455	11.49	0.0009	***	2.997
OM	10.0	41.696	10.94	0.0011	**	2.904
CP	10.0	58.231	14.62	0.0004	***	3.082
NDF	10.0	20.371	0.16	0.955	ns	17.556
ADF	10.0	10.836	9.42	0.002	**	2.007

*DF: error degree of freedom; MS: mean square of treatments; Pr: probability; CV: coefficient of variance; SL: significant level; ns: non-significant; ADF: acid detergent fiber; CP: crude protein; NDF: neutral detergent fiber; OM: organic matter; DM: dry matter*

**Appendix Table 2. Summary of ANOVA for total dry matter and nutrient intake of experimental sheep**

Intake (gd <sup>-1</sup> )	DF	MS	F-value	Pr >F	SL	CV (%)
Hay DM intake	20.0	5103.94	13.91	<.0001	***	5.09
Total DM	20.0	5103.94	13.91	<.0001	***	2.833
Total OM	20.0	4435.838	5.18	0.0050	**	4.298
Total CP	20.0	29.657	6.10	0.0022	**	2.662
Total NDF	20.0	9998.526	11.70	<.0001	***	6.089
Total ADF	20.0	14556.837	18.39	<.0001	***	8.429
ME (MJ/d)	10.0	1.0062	7.31	0.0051	**	4.894

*DF: error degree of freedom; DM: dry matter; MS: mean square of treatments; Pr: probability; CV: coefficient of variance; SL: significant level; ADF: acid detergent fiber; CP: crude protein; NDF: neutral detergent fiber; OM: organic matter; ME: metabolizable energy*

### List of Figures in the Appendix:



**Appendix Fig. 1. Lambs arranged in the pen**



**Appendix Fig. 2. Technique/procedure for collecting excrement**

**Disclaimer/Publisher's Note:** The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of the publisher and/or the editor(s). This publisher and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.

© Copyright (2024): Author(s). The licensee is the journal publisher. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

*Peer-review history:*  
The peer review history for this paper can be accessed here:  
<https://www.sdiarticle5.com/review-history/123661>