



Genetic Variability Studies in F₂ Populations of Ridge Gourd [*Luffa acutangula* L. (Roxb.)] for Growth, Yield and Quality Traits

Kavya^{a++*}, H. P. Hadimani^{a#},
Shashikanth Evoor^{a†}, Satish D^{b†} and Venkateshalu^{c‡}

^a Department of Vegetable Science, College of Horticulture, UHS, Bagalkot, India.

^b Department of Genetics and Plant Breeding, College of Horticulture, UHS, Bagalkot, India.

^c Department of Entomology, College of Horticulture, Kolar, UHS, Bagalkot, India.

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

DOI: <https://doi.org/10.9734/jabb/2024/v27i101582>

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/125458>

Original Research Article

Received: 15/08/2024

Accepted: 17/10/2024

Published: 26/10/2024

ABSTRACT

An experiment was conducted to study, genetic variability in ridge gourd during *rabi* 2024 at College of Horticulture, Bagalkot. The observations were recorded on various growth, yield and quality contributing characters of population-I (ArkaPrasan x Sirsi Local-2) and population-II (Jaipur Long x KLR-5). The analysis of variance indicated the prevalence of sufficient genetic variation among the

⁺⁺ M.sc. (Hort.) Vegetable Science;

[#]Assistant Professor;

[†] Associate Professor;

[‡]Professor;

*Corresponding author: E-mail: kummankavya@gmail.com;

Cite as: Kavya, H. P. Hadimani, Shashikanth Evoor, Satish D, and Venkateshalu. 2024. "Genetic Variability Studies in F₂ Populations of Ridge Gourd [*Luffa Acutangula* L. (Roxb.)] for Growth, Yield and Quality Traits". *Journal of Advances in Biology & Biotechnology* 27 (10):1578-86. <https://doi.org/10.9734/jabb/2024/v27i101582>.

genotypes from all the characters studied. The high phenotypic coefficient of variation (PCV) and genotypic coefficient of variation (GCV) were observed for node at first male flower appears, node at first female appears, TSS and crude fibre in population-I. Whereas, the traits like vine length, number of primary branches per vine, average fruit weight, fruit yield per vine and crude fibre in population-II. High heritability coupled with high genetic advance were observed for node at first male flower appears, node at first female flower appears, fruit length, fruit diameter, average fruit weight, fruit yield per vine, number of fruits per vine, number of seeds per fruit, fruit tenderness, TSS, crude fibre and vitamin C in the both F₂ populations indicating these characters are governed by additive gene action. Hence, direct selection may be followed for the improvement of ridge gourd for these characters.

Keywords: Ridge gourd; variability; heritability and genetic advance.

1. INTRODUCTION

“Ridge gourd *Luffa acutangula* L. (Roxb.) with a diploid chromosome number of $2n=2x=26$, belongs to the Cucurbitaceae family and is an important vegetable crop cultivated in tropical and subtropical regions worldwide” [1]. It is a nutrient-rich fruit, containing about 92.5 grams of moisture and providing essential nutrients like calcium and vitamin C, along with only 17 kilocalories per 100 grams. Ayurvedic literature highlights its health benefits, including enhancing the immune system, reducing blood sugar levels, and serving as a diuretic. The seeds of ridge gourd contain cucurbitacin, which has purgative and anthelmintic properties, along with ribosome inactivating proteins (RIPs) that show potential in medicinal applications, such as inhibiting tumor growth and exhibiting antiviral effects. Overall, it is recognized for its significant nutritional and therapeutic value [2]. Various genotypes and cultivars of ridge gourd are grown across different areas of the country, each exhibiting diverse characteristics. This variability among genotypes can result from geographical differences or natural cross-pollination. A thorough understanding of the existing genetic variability in a crop is crucial before initiating any breeding program [3]. The phenotypic traits of plants are primarily influenced by their genetic makeup and the environmental conditions they encounter, as well as the interaction between genotypes and the environment. “Plant genotype is determined by both heritable and non-heritable factors, making it essential to differentiate observed phenotypic variability into heritable and non-heritable components. In crop improvement, the genetic variation is important as this component is transmitted from one generation to another generation. The effectiveness of selection for any trait depends not only on the amount of phenotypic and genotypic variability, but also on the magnitude of heritability. The

magnitude of heritability indicates the effectiveness with which selection of genotypes can be made based on phenotypic performance” [4]. Thus the exploitation of variability is of great importance and is pre-requisite for any crop improvement. Considering the above facts the F₂ population has large variability, it provides ample scope for screening or selecting of best plants. Therefore, the present investigation was undertaken.

2. MATERIALS AND METHODS

The experiment was carried out at College of Horticulture, University of Horticultural Sciences, Bagalkot. It is situated in Northern dry zone of Karnataka state at 16° 46' North latitude and 74° 59' East longitude and an altitude of 533.0 meters above the mean sea level. Crop raised during rabi season in the months of January-May, 2024. In order to develop the experimental material required for the variability studies in F₂ generation. Population-I (ArkaPrasan x Sirsi Local-2), population-II (Jaipur Long x KLR-5) these two crosses hybrids were used for Selfing in order to generate F₂ populations. These F₂ seeds were sown to study the variability on 14th Jan 2024 at field of Vegetable Science, College of Horticulture, Bagalkot. Seeds were treated with butter milk for 24 hours for better germination. Treated seeds were sown in trays which were filled with coco peat and seedlings were transplanted at 16 days from sowing in the trays, were planted in the main field. The seedling were placed on a raised bed with a spacing of 2m x 1m, mulching was implemented utilizing a 30-micron black polythene mulch sheet and watering was done by using driplines which were placed on bed before planting. The experiment was conducted by using augmented design. The crop received timely management practices as per recommended package of practices [5]. The crop

was maintained properly till last harvest and observations on growth as well as yield contributing characters was noted on F₂ progenies along with parents. From each cross 200 plants were studied and taken data from all the plants. Two hundred plants each from two hybrids were maintained in this experiment. Genotypic and phenotypic coefficient of variation were calculated as per the formula suggested by Burton and vane [6]. Heritability and expected genetic advance were calculated as per formula given by Johnson et al., [4].

The observations on vine length at harvest time (m), the number of primary branches at 45 DAP, days to first male flowering, days to first female flowering, the node of first female flowering, the node of first male flowering, sex ratio (male to female), days to first harvest and days to last harvest, number of fruits per vine, fruit length (cm), fruit weight (g), fruit diameter (mm), fruit yield per vine (kg), number of seeds per fruit, fruit tenderness (N), TSS (°Brix), crude fibre(%) and vitamin C (mg/ 100 g) were recorded in both populations on individual plants. Tenderness of fruits were estimated by piercing the fruits using penetrometer.

3. RESULTS AND DISCUSSION

“The results in the current study showed that the phenotypic coefficient of variation (PCV) values are higher than the genotypic coefficient of variation (GCV), suggesting that environmental factors have a significant impact on the performance of the F₂ population. The findings related to genetic variability, heritability and genetic advance for various traits are discussed in this study are presented in Table 1 and Table 2. Higher estimates of genotypic coefficient of variation (GCV) and phenotypic coefficient of variation (PCV) were observed for node at first male flower appearance, node at first female appearance, TSS and crude fibre were found in population-I. Whereas, the traits like vine length, number of primary branches per vine, average fruit weight, fruit yield per vine and crude fibre were observed in population-II. The existence of a broad genetic base within the population is shown by high GCV and PCV values with small differences. This suggests consistent genotype expression for these traits, allowing for the potential for these qualities to be improved through simple selection. The similar results were also observed” by Choudhary and Kumar [7], Saklesh [8], Bhargava et al. [9] in ridge gourd.

“Moderate amount of GCV and PCV were observed for the traits like vine length, fruit length, fruit diameter, average fruit weight, number of fruits per vine, number of seeds per fruit, fruit tenderness and vitamin C in population-I. In population-II it was observed for traits like node at first female flower appearance, node at first male flower appearance, fruit diameter, number of fruits per plant, number of seeds per plant, fruit tenderness and vitamin C. As this character had moderate PCV and GCV representing that the trait is controlled by both additive and non-additive gene actions. Hence, recurrent selection can be practiced for improvement. These observations are in line with the findings of” Varalakshmi et al. [10], Singh et al. [11] and Panda et al. [12] in ridge gourd.

“Moderate GCV coupled with high PCV was observed for number of primary branches per vine and fruit yield per vine in population-I and for the traits like fruit length and TSS it was observed in population-II. Hence, the presence of moderate GCV and high PCV for a trait in a genetic study could indicate diverse underlying genetic factors like polygenic inheritance, epistasis, presence of genetic mutations or allelic variations, etc. contributing to the observed variability. These results are in accordance with” Dubey et al. [13] and Tiwari et al. [14], Choudhary et al. [7], Koppad et al. [15] and Panda et al. [12].

“Lower GCV and moderate PCV was observed for number of male flowers, number of female flowers and sex ratio in population-I and for the traits like number of male flowers and number of female flowers it was observed in population-II, which indicates the presence of moderate amount of variability for these traits and little influence of environment on these traits. These observations are in line with the findings” of Manoj et al. [16], Harshitha et al. [17] and Thulasiram et al. [18].

“Low magnitude of GCV and PCV were recorded for days to first male flower, days to first female flower, days to first fruit harvest and days to last fruit harvest in population-I. In population-II it was observed for days to first male flower, days to first female flower, days to first fruit harvest, days to last fruit harvest and sex ratio. In accordance” with Choudhary et al. [7], Koppad et al. [15], Singh et al. [19], Manoj et al. [16], Methela et al. [20], Harshitha et al. [17], Kannan et al. [21], Akhila and Singh [22], Sravani et al. [23], Durga et al. [24] and Thulasiram et al. [18].

Table 1. Genetic variability estimates for various traits in the F₂ generation of population-I (ArkaPrasan x Sirsi Local-2)

Sl. No	Traits	Mean	Range	PCV (%)	GCV (%)	h ² bs (%)	GAM (%)
1	Vine length at harvest time (m)	3.48	1.82- 4.59	14.29	11.10	60.36	17.79
2	Number of primary branches per vine (45DAS)	3.01	1.00-5.00	22.01	18.63	71.68	32.50
3	Days to first female flower	39.40	36.00-49.00	6.66	5.81	76.03	10.45
4	Days to first male flower	35.66	32.00-46.00	7.77	7.28	87.70	14.06
5	Node at first male flower appearance	5.19	3.00-14.00	38.90	36.16	86.42	69.34
6	Node at first female flower appearance	9.20	5.00-19.00	25.08	23.49	87.72	45.39
7	Number of male flowers	523.38	346.00-656.00	11.51	9.96	74.88	17.77
8	Number of female flowers	21.97	14.00-28.00	11.38	7.49	43.30	10.17
9	Sex ratio (M: F)	23.96	18.78-34.63	11.05	8.24	55.69	12.69
10	Days to first fruit harvest	46.19	44.00-53.00	3.98	3.23	65.83	5.41
11	Days to last fruit harvest	105.81	78.00-125.00	8.25	6.83	68.56	11.67
12	Fruit length (cm)	27.76	15.00- 41.50	14.55	12.70	76.20	22.88
13	Fruit diameter (mm)	35.64	20.90-49.20	12.19	10.98	81.07	20.39
14	Average fruit weight (g)	163.84	78.00-250.70	18.72	16.87	81.23	31.37
15	Fruit yield per vine(kg)	2.31	0.96-3.40	22.73	19.72	75.21	35.27
16	Number of fruits per vine	14.13	6.00-20.00	16.09	13.39	69.22	22.98
17	Number of seeds per fruit	119.92	72.00-180.00	16.61	13.08	62.04	21.26
18	Fruit tenderness (N)	6.28	3.68-8.54	19.98	18.49	72.66	33.56
19	TSS (° B)	3.29	2.10-8.21	24.12	22.04	83.50	41.54
20	Crude fibre (%)	0.62	0.22-1.61	30.89	29.30	89.99	57.34
21	Vitamin- C (mg/100g)	10.65	0.28-13.74	12.01	11.47	91.21	22.60

GCV- Genotypic coefficient of variation; h²bs- Heritability (Broad sense);

PCV- Phenotypic coefficient of variation; GAM- Genetic advance over mean; S. Em- Standard Error deviation from mean

Table 2. Genetic variability estimates for various traits in the F₂ generation of population-II (Jaipur Long x KLR-5)

Sl. No	Traits	Mean	Range	PCV (%)	GCV (%)	h ² bs (%)	GAM (%)
1	Vine length at harvest (m)	3.03	1.62- 4.33	22.83	21.38	87.69	41.30
2	Number of primary branches per vine (45DAS)	2.75	1.00- 5.00	37.64	36.39	93.42	72.55
3	Days to first female flower	40.01	37.00- 51.00	8.83	6.45	81.42	11.43
4	Days to first male flower	36.14	34.00-39.00	6.53	4.73	60.72	10.35
5	Node at first male flower appearance	5.78	4.00-9.00	19.12	17.48	77.17	32.79
6	Node at first female flower appearance	10.43	5.00-14.00	14.63	11.86	71.63	21.13
7	Number of male flowers	520.18	346.00-675.00	11.35	9.42	68.93	16.14
8	Number of female flowers	21.17	14.00-28.00	11.23	9.38	69.74	16.15
9	Sex ratio	24.70	19.23-31.00	9.99	8.90	79.46	16.38
10	Days to first fruit harvest	46.99	44.00-51.00	5.64	3.80	46.56	5.53
11	Days to last fruit harvest	106.97	85.00-120.00	9.35	7.27	51.34	10.79
12	Fruit length (cm)	25.24	12.78-38.90	20.11	19.47	93.76	38.90
13	Fruit diameter (cm)	36.56	21.29-55.30	17.70	17.19	94.24	34.42
14	Average fruit weight (g)	129.42	59.40-226.00	23.49	21.71	85.44	41.40
15	Fruit yield per vine(kg)	1.79	0.66-2.71	26.08	22.95	77.41	41.65
16	Number of fruits per vine	13.85	9.00-22.00	15.94	13.58	72.50	23.85
17	Number of seeds per fruit	105.49	61.00-186.00	17.40	13.44	60.66	21.42
18	Fruit tenderness (N)	6.12	3.45-9.20	15.70	12.79	66.37	21.50
19	TSS (° B)	3.21	1.90-5.40	20.59	16.75	66.18	28.11
20	Crude fibre (%)	0.48	0.09-1.23	38.40	36.87	92.18	33.03
21	Vitamin- C (mg/100g)	10.55	7.17-13.70	10.47	10.15	93.93	20.29

GCV- Genotypic coefficient of variation; h²bs- Heritability (Broad sense)

PCV- Phenotypic coefficient of variation; GAM- Genetic advance over mean; S. Em- Standard Error

This indicates the low variability for these parameters in the germplasm and preponderance of non-additive gene action. This offers very limited opportunity for improving these traits by selection. This suggests the narrow genetic base and hence the variability has to be generated for these characters through hybridization to recover transgressive segregants or through mutation breeding.

“High estimates of heritability coupled with high values of GAM were observed for the characters viz., number of primary branches, node at first male flower appearance, node at first female flower appearance, fruit length, fruit diameter, average fruit weight, number of fruits per vine, fruit yield per vine, number of seeds per fruit, TSS, crude fibre, fruit tenderness and vitamin C in population-I. Whereas, the traits like vein length, number of primary branches, node at first male flower, node at first female flower appearance, fruit length, fruit diameter, average fruit weight, fruit yield per vine, number of fruits per vine, number of seeds per fruit, fruit tenderness, TSS, crude fibre and vitamin C in population-II. This indicates predominance of additive components for these traits and hence direct selection would be more effective in improving these traits. These results are accordance with matching conclusions were confirmed” by Singh et al. [19], Choudhary et al. [7], Dubey et al. [13], Varalakshmi et al. [10], Harshitha et al.[17] and Thulasiram et al. [18].

High heritability coupled with moderate GAM were observed for the traits viz., vine length, days to first male flower, days to first female flower, days to last harvest and number of male flowers in population-I. Whereas, in population-II it was observed for days to first female flower appearance, days to first male flower appearance, number of male flowers, number of female flowers and sex ratio. This indicates predominance of additive components deviation from mean for these traits and hence direct selection would be more effective in improving these traits. These results are accordance with findings of Manoj et al. [16], Kannan et al. [21], Akhila and Singh [22] and Thulasiram et al. [18].

Moderate heritability and GAM were observed for number of female flowers and sex ratio in population-I. Whereas, the trait days to last harvest in population-II, which indicated limited scope for improvement of these characters through direct selection. These results are

accordance with Harshitha et al. [17], Singh et al. [19] and Panda et al. [12]. High heritability coupled with low GAM were observed for days to first fruit harvest in population-I and population-II, which indicated the effect of the environment on the expression of particular traits. A high heritability with low genetic advance may also indicate non-additive gene action and selection for such trait not be effective. These results are in accordance with Kumari et al. [25], Akhila and Singh [22] and Thulasiram et al. [18].

4. CONCLUSION

The analysis of F₂ data in both populations indicate the presence of good amount of genetic variation for most of characters. High GCV and PCV were observed for node at first male flower appears, node at first female appears, TSS and crude fibre in population-I. Whereas, the traits like vine length, number of primary branches per vine, average fruit weight, fruit yield per vine and crude fibre in population-II. High heritability coupled with high genetic advance as per cent over mean were observed for node at first male flower appears, node at first female flower appears, fruit length, fruit diameter, average fruit weight, fruit yield per vine, number of fruits per vine, number of seeds per fruit, fruit tenderness, TSS, crude fibre and vitamin C in the both F₂ populations. Additive inheritance is predominant in these characters. This indicates predominance of additive components for these traits and hence direct selection would be more effective in improving these traits.

DISCLAIMER (ARTIFICIAL INTELLIGENCE)

Author(s) hereby declare that NO generative AI technologies such as Large Language Models (ChatGPT, COPILOT, etc) and text-to-image generators have been used during writing or editing of this manuscript.

ACKNOWLEDGEMENTS

The author extends sincere gratitude to the Department of Vegetable Science, College of Horticulture, Bagalkot, for providing financial support and necessary facilities during the research period.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

- Bose TK, Som MG. Vegetable crops in India. Nayapokash, Bidhausarani, Calcuta. India. 1986;107-114.
- Manikandaselvi S, Vadivel V, Brindha P. Review on *Luffa acutangula* L. Ethnobotany, phytochemistry, nutritional value and pharmacological properties. Int. J. Curr. Pharm. Rev. Res. 2016;7(3):151-5.
- Singh BD, Principles of Genetics, Kalyani Publishers. 1992;78-100.
- Johnson HW, Robinson HF, Comstock RE. Genotypic and phenotypic correlations in soyabean and their implications in selection. J. Agron. 1955;37:477-483.
- Anonymous. Integrated handbook of horticulture (Kannada), Univ. Hort. Sci. 2022;103-104.
- Butron GW, De Vane EH. Estimating heritability in tall-fescue (*Festuca arundinacea*) from replicated clonal material. Agron. J. 1953;45:478-481.
- Choudhary BR, Kumar S. Genetic analysis in ridge gourd [*Luffa acutangula* (Roxb.) L.] under hot arid conditions. Indian J. Genet. Plant Breed. 2011;28(8):156-168.
- Saklesh. Genetic variability studies in ridge gourd. M.Sc. (Agri.) Thesis, Orissa Univ. Agric. and Tech., Bhubaneshwar (India); 2016.
- Bhargava AK, Singh VB, Kumar P, Meena RK. Efficiency of selection based on genetic variability in ridge gourd (*Luffa acutangula* L. (Roxb.)). J. Pharmacogn. Phytochem. 2017;6(4):1651-1655.
- Varalakshmi B, Pitchaimuthu M, Rao ES, Manjunath KSS, Swathi SH. Genetic variability, correlation and path analysis in ridge gourd (*Luffa acutangula* L.). J. Hort. Sci. 2015;10(2):21-26.
- Singh RP, Mohan J, Dharmendra S. Studies on genetic variability and heritability in ridge gourd (*Luffa acutangula* L.). Agric. Sci. Dig. 2020;22(4):279-280.
- Panda M, Mohanty A, Sarkar S, Sahu GC, Tripathy P, Das SK, Patnaik A. Variability studies in ridge gourd (*Luffa acutangula* (L.) Roxb.). J. Pharm. Innov. 2022;11(4):1716-1719.
- Dubey RK, Singh V, Upadhyay G. Genetic variability and inter relationship among some ridge gourd (*Luffa acutangula* L.) accessions under foot hills of Arunachal Pradesh. Prog. Hort. 2013;45(1):191-197.
- Tiwari NK, Pandey AK, Sing UN, Singh VB. Genetic variability, heritability in narrow sense and genetic advance per cent of mean in bitter gourd (*Momordica charantia* L.). J. Pharmacogn. Phytochem. 2018;7(2):2608-2610.
- Koppad SB, Chavan M, Hallur RH. Character association studies and path coefficient analysis for yield and yield attributing traits in ridge gourd (*Luffa acutangula* (L.) Roxb.). Electronic J. Plant Breed. 2016;7(2):275-281.
- Manoj YB, Lakshmana D, Ganapathi M, Chandana BC. Studies on character association and genetic variability for important traits in ridge gourd (*Luffa acutangula* L.). Green Farming Int. J. 2018;9(2):244-247.
- Harshitha S, Sood M, Indiresk KM. Variability and heritability studies for horticultural traits in ridge gourd (*Luffa acutangula* (L.) Roxb.). Int. J. bio-resour. Stress Manag. 2019;10(4):335-339.
- Thulasiram LB, Ranpise SA, Bhalekar MN. Variability studies in ridge gourd (*Luffa acutangula* L. Roxb.). Int. J. Veg. Sci. 2022;2(5):167-177.
- Singh V, Rana DK, Shah KN. Genetic variability, heritability and genetic advance in some strains of bitter gourd (*Momordica charantia* L.) under subtropical conditions of garhwahimalaya. Plant Arch. 2017;17(1):564-568.
- Methela NJ, Islam MS, Mitu N, Latif MA. Mean performance and heritability studies of some ridge gourd (*Luffa acutangula* L.) genotypes. Research in Agriculture Livestock and Fisheries., 2019;6(1):45-55.
- Kannan A, Rajamanickam C, Krishamoorthy V, Arunachalam P. Genetic variability, correlation and path analysis in F₄ generation of (*Luffa acutangula* (Roxb) L.). Int. J. Chem. Stud. 2019;7(3):208-213.
- Akhila K, Singh D. Genetic variability in ridge gourd (*Luffa acutangula* (L.) Roxb.). Int.J. Curr. Microbiol. Appl. Sci. 2020;9(10):2774-2783.
- Sravani Y, Rekha GK, Ramana CV, Naidu LN, Suneetha DS. Studies on genetic variability, heritability and genetic advance in F₂ generation of ridge gourd. J. Pharm. Innov. 2021;10(7):927-930.
- Durga PM, Chinthalapudi, Kranthi Rekha G, Usha Kumari K, Uma Jyothi K, Narasimharao S. Variability studies in F₃ population of ridge gourd (*Luffa acutangula*) for yield and yield attributing traits. Pharma innov. 2012;10(7):612-615.

25. Kumari M, Kumar J, Kumari A, Singh VK, Rani N, Kumar A. Genetic variability, correlation and path coefficient analysis for yield and yield attributing traits in bitter gourd (*Momordica charantia* L.). Curr. J. Appl. Sci. Technol. 2018;31(4):1-8.

Supplementary Material 1. Mean performance of parents and hybrids for growth and yield parameters

Parents	VL	PBPT	DFP	DFM	NMF	NFF	DFHT	DLHT	SR	FL	FD	AVGWT	NFPV	NSF	FYPV	FT	TSS	CF	Vit C
ArkaPrasan (AP)	3.55	3.82	37.79	34.98	5.00	10.00	45.14	109.71	21.30	30.78	36.94	152.29	16.29	194.14	2.47	6.39	2.98	0.60	12.09
Sirsi Local-2 (SL)	3.10	3.29	40.24	35.92	5.14	10.14	47.29	108.43	22.48	26.66	32.14	132.43	15.29	165.14	2.02	6.64	2.77	0.74	12.47
KLR-5	3.68	3.19	41.16	38.02	6.00	11.29	46.86	105.71	23.17	28.89	33.35	117.42	16.71	122.57	1.96	6.40	2.52	0.62	12.41
Jaipur Long (JP)	3.60	3.13	40.91	36.53	6.71	11.29	45.86	102.86	21.79	19.40	33.04	125.93	15.57	122.43	1.96	5.60	3.60	0.59	10.98
F ₁ (AP× SL)	3.44	3.02	39.84	36.01	5.43	10.14	46.29	106.57	22.22	26.25	33.25	153.71	16.43	123.14	2.50	5.68	3.62	0.70	11.02
F ₁ (JP× KLR-5)	3.70	2.72	40.57	36.29	8.57	13.86	47.43	107.71	22.43	29.39	34.79	132.20	17.00	135.71	2.21	6.27	4.05	0.30	13.01

Disclaimer/Publisher's Note: The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of the publisher and/or the editor(s). This publisher and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.

© Copyright (2024): Author(s). The licensee is the journal publisher. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:
 The peer review history for this paper can be accessed here:
<https://www.sdiarticle5.com/review-history/125458>