



Study on Crop-weather Calendar of Wheat for Eastern Plain Zone of Uttar Pradesh

Anushka Pandey^{a*}, A. K. Singh^a, A. N. Mishra^a and S. R. Mishra^a

^a *Department of Agricultural Meteorology, A.N.D.U.A.T. Kumarganj, Ayodhya, UP. 224229, India.*

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

DOI: 10.9734/IJECC/2022/v12i1130965

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/82526>

Original Research Article

Received 01 May 2022

Accepted 05 July 2022

Published 09 July 2022

ABSTRACT

An investigation entitled "Study on Crop-weather calendar of wheat crop for eastern plain zone of Uttar Pradesh" was carried out at the Department of Agricultural Meteorology, ANDUA&T, Kumarganj, Ayodhya. Crop weather calendar for wheat crop has been prepared for district Sultanpur through the collection of historical weather data of last 20 years (2000- 2020). Crop weather calendar of wheat was formulated by combining the weekly climatic averages and phenological calendar for the crop along with optimum weather criteria needed at different phenological stages of the crop. Climatic normal for wheat crop has been taken from 46th Standard meteorological week to 14th Standard meteorological week (from sowing to harvesting). From the crop weather calendar of wheat crop it was observed that for during seed emergence, T max 25-27^oC, T min 11.5-13^oC, RHm 92.5-92.8% and RHe 42-49%, at CRI, T max 24-26^oC and T min 7.5-11^oC, and during milking stage of the crop, T max 30-33^oC, T min 12.3-14^oC, RHm 88-91%, RHe 43% were found conducive for better yield at district Sultanpur. Conducive weather conditions for infestation of army worm were T max 20-29^oC and T min 14-8^oC, RHm 90-94% and RHe 49-65% and rainfall of 1.5- 11 mm from met. no. week 46th to 4th met. week and for leaf blight T max 20-28.5^oC, T min 12-17^oC, RHm 88-91% and RHe 42-52% from 1st to 10th met. week at booting and milking stage of the crop.

Keywords: *Crop-weather calendar; wheat; climatic normal.*

1. INTRODUCTION

Wheat (*Triticum aestivum* L.) belongs to the *poaceae* family. Wheat production is estimated to increase to a record 109.52 million tons in 2020-21 with an average national productivity of 3424 kg/ha. from 107.86 million tons in the previous year, while the output of coarse cereals is likely to increase to 51.15 million tons from 47.75 million tons. (II Advance Estimates of 2020-21, DAC&FW). Uttar Pradesh has been the largest producer state during 2019-2020 with production of 32.59 million tones followed by MP 19.61 million tons and Punjab 17.57 million tons. The wheat production is distributed in three agroclimatic zones of UP viz. western Uttar Pradesh (3.29 million ha) ,eastern uttar pradesh (5.24 million ha) and central Uttar Pradesh (0.68 million ha). (icar.org.in).

It is sown in mid-October to mid- November and harvested in March- April. Wheat is widely adapted crop. It's grown from temperate, irrigated to dry and high rainfall areas and from warm, humid to dry, cold environments. Wheat is a cool season crop, hence cool weather during vegetative development and warm weather for maturity is deemed ideal for wheat. During the heading and flowering phases, exceedingly high or low temperatures and drought are detrimental to wheat. Wheat production is considerably reduced by temperatures above the optimum. According to Porter and Gawith [1], the optimum temperature for wheat is around 20°C, ranging between 17°C and 23°C; plants stop to grow below 0°C or above 37°C, and die at around - 17°C or 47.5°C. Increased daytime and nighttime temperatures decrease wheat yields in controlled environments [2].

Detailed information of every important crop on their dates of sowing dates of commencement and duration of general cultural operations, vital periods in their life cycle and their most usual weather demands have in India been presented in a very pictorial form called the Crop Weather Calendar [3].

Crop weather calendar may be a comprehensive guide for researcher and farmers. It is a resource that provides information on average weather of every week, planting, sowing and harvesting periods of locally adopted crops in a specific agro-ecological zone [4-6].

It also provides information on the sowing rates of seed and planting material and the so the most agricultural practices. This tool supports

farmers and agriculture extensionists in taking appropriate decisions on crops and their sowing period, respecting the agro-ecological dimension. It also provides a strong base for emergency/contingency planning of the rehabilitation of farming systems after disasters [7].

The objectives of this study are to find out the suitable climatic normal and the phenophase wise congenial weather for the better yield of wheat crop and to present it on a single sheet of paper, so it can be used by forecasters to issue the appropriate agro-advisory.

2. MATERIALS AND METHODS

Eastern plain zone UP covers the 10 districts namely Barabanki, Faizabad, Sultanpur, Jaunpur, Azamgarh, Mau, Ballia, Ghazipur, Varanasi and Sant Ravidas Nagar. Rainfall is adequate with a normal of 1,025 mm. The climate is dry sub-humid to moist sub-humid. Over 70% of the land is cultivated and more than 80% of the cultivated area is irrigated.

Weather data for last twenty years (2000-2020) were collected from Department of Agricultural Meteorology, ANDUA&T, Kumarganj, Ayodhya and IMD. Weekly climatic normal for standard meteorological weeks (1st - 52nd) for this location were computed (Table 1). These meteorological data sets were arranged in weekly format for cropping season from the month of sowing to the month of harvest of the crop. The collected weather data of Sultanpur district were arranged and calculated to find out average, summation, maximum and minimum value which was used for further analysis.

Crop weather calendar of wheat crop for district Sultanpur was formulated by combining the weekly climatic averages and phenological calendar for the crop along with optimum weather criteria needed at different phenological stages of the crop. Processing of data worked on the basis of standard date of sowing recommended for given crops and the standard average duration of crop. The collected weather data was tabulated to find out average value of weather parameters viz. Tmin, Tmax, Rainfall, morning RH (RHm), evening RH (RHe) and Evaporation for further analysis. The range of different meteorological parameters for the higher production of wheat crop at district Sultanpur was worked out from the actual meteorological data of high productivity crop year which was collected form the ATIC of ANDUA&T, Kumarganj Ayodhya Table 2.

Table 1. Climatic normal for weather parameters (2000-2020)

Week/ weather parameter	T min(°C)	T max(°C)	RHm (%)	RHe(%)	Rainfall (mm)	Evp (mm/day)	Bss (hrs.)
1	6.7	19.1	89.8	63.8	4.7	12.0	4.49
2	6.1	19.3	85.9	61.1	1.1	13.5	4.72
3	7.2	20.8	86.6	57.6	5.2	14.1	5.71
4	7.0	21.3	88.2	56.3	5.3	15.4	7.28
5	7.7	22.8	87.7	53.2	1.3	18.2	6.73
6	8.6	24.5	90.0	52.7	5.7	20.6	7.18
7	9.6	23.5	86.1	51.1	3.5	21.4	8.7
8	10.8	27.0	86.2	46.2	5.5	25.1	8.49
9	11.9	27.8	83.6	43.8	3.6	26.4	8.32
10	12.2	29.5	79.9	42.1	4.2	27.2	8.56
11	13.5	30.6	78.6	40.4	2.5	30.3	8.39
12	14.9	33.2	74.1	34.9	0.4	34.5	8.38
13	16.1	34.0	71.7	30.6	0.4	36.6	8.7
14	17.0	36.3	68.8	31.1	0.7	40.8	8.81
15	19.2	36.3	62.5	28.8	0.8	48.3	8.81
16	21.1	37.9	66.6	32.1	1.4	47.0	9
17	21.7	40.1	62.9	31.4	3.5	50.5	9.15
18	23.0	37.4	69.2	32.8	5.1	48.7	9.34
19	23.8	39.0	67.0	33.5	5.1	52.6	9.3
20	24.0	39.1	69.4	34.7	3.7	54.0	9.18
21	25.6	38.6	70.6	36.7	5.5	52.4	8.41
22	25.4	38.7	69.2	40.2	10.9	51.7	8.6
23	25.9	38.5	73.2	43.8	25.1	51.4	8.13
24	26.5	37.7	72.8	48.1	22.7	47.5	7.24
25	26.7	35.7	80.3	57.7	38.3	41.9	5.69
26	26.7	35.4	82.2	59.6	66.2	37.0	4.57
27	26.5	33.5	85.0	70.2	68.4	32.4	4.89
28	25.8	33.1	87.8	71.3	57.7	36.0	4.11
29	26.0	32.9	87.9	72.5	71.0	31.1	4.85
30	26.3	32.8	88.6	72.4	45.1	29.3	5.68
31	26.2	32.0	87.9	71.7	44.4	29.3	4.92
32	26.4	33.3	85.7	70.6	40.8	32.6	4.95
33	26.1	33.1	90.4	74.7	67.0	27.4	4.38
34	26.2	32.4	89.6	73.4	51.7	31.0	4.45
35	26.0	29.5	89.2	70.1	30.8	33.6	5.42
36	25.8	32.9	86.9	70.2	37.4	31.6	4.59
37	25.6	32.9	89.4	70.1	36.5	32.5	5.06
38	25.0	32.7	89.1	69.7	35.9	31.8	6.07
39	23.8	32.6	88.3	69.1	39.7	30.5	6.52
40	22.9	32.6	86.2	64.7	11.3	32.8	7.17
41	21.0	32.9	85.3	59.8	6.7	32.2	7.5
42	19.2	32.1	86.8	52.6	5.2	31.1	8.19
43	16.9	31.3	87.1	50.9	0.5	28.3	7.69
44	15.4	30.4	87.5	48.3	0.0	27.0	8.26
45	13.9	29.9	89.0	49.6	0.3	25.8	7.89
46	13.2	28.9	88.8	49.2	0.6	25.3	7.37
47	11.6	27.5	85.0	46.5	0.0	23.0	7.14
48	10.5	27.2	87.2	46.7	0.0	21.3	6.8
49	8.9	25.4	87.9	47.9	0.0	30.9	6.85
50	8.7	24.0	88.5	53.4	2.0	19.7	6.59
51	7.3	22.2	89.9	56.7	0.0	22.2	5.88
52	6.6	22.2	91.2	58.3	1.4	30.2	5.77

Table 2. Highest Productivity Year for wheat at Sultanpur

Year	2015	2016	2018	2020
Productivity of wheat qha ⁻¹	286.49	261.03	354.71	322.66




Fig. 1. Structure of crop weather calendar

The data on weather condition favorable for incidence of pest and nature of weather warnings were collected from the Entomology Department of ANDU&T Kumarganj Ayodhya. The investigation is based on data collection of pest investigation on wheat and weather and averaged over last 5 years.

2.1 Structure of Crop-weather Calendar

Structure of crop weather calendar consists of main three parts viz: upper part, which contains the location specific weather data for the crop growing season, middle part which shows the typical life history of the crop in the form of diagram. Important “growth phases” related to the crop like sowing, germination/ emergence, transplanting (in the case of rice), vegetative growth, flowering, grain formation and maturity period etc. are indicated. In addition, suited to the above acquaintance, the middle part of the calendars revealed the compatible meteorological condition for the crop (stage-wise or whole crop growth period) which will lead towards high yield of the crop. In the bottom part favorable weather conditions for the incidence of pest and diseases are reported (Fig. 1).

3. RESULTS AND DISCUSSION

3.1 Crop Weather Calendar for Wheat (District Sultanpur) Climatic normal for Wheat at Sultanpur

Climatic normal for wheat crop has been taken form Standard meteorological week 46th to 14th met. week (from sowing to harvesting). The highest normal rainfall (5.43 mm) was found during 2nd met. week followed by 5.33mm during 3rd met. week. The highest normal T max (36^oC) and T min (16.9^oC) were found during 14th week. Highest morning RH (RHm) was during 52nd week (91%) and minimum evening RH (RHe) found during 13th week (30%) (Fig. 2).

3.2 Phenophase-wise Weather for Better Yield for Wheat at District Sultanpur

Wheat seed had taken 6-7 days to emerge. During seed emergence, T max 25-27^oC, T min 11.5-13^oC, morning RH (RHm) 92.5-92.8% and evening RH (RHe) 42-49% were found conducive for better yield. The optimum temperature range for the early growth stages of wheat is lower than the threshold for the later growth stages because

wheat production can be considerably reduced by temperatures above the optimum. Narayanan et al. [8] also reported that increased daytime and nighttime temperatures decrease wheat yields in controlled environments.

At CRI stage of the crop, T max and T min 24-26°C and 7.5-11°C, respectively, and at tillering stage T max 15-23°C, T min 5-3.5°C, morning RH (RHm) 94-97%, evening RH (RHe) 47-48% and BSS >4 hrs/day were found conducive for district Sultanpur. During milking stage of the crop, T max 30-33°C, T min 12.3-14°C, morning RH (RHm) 88-91%, RHe 43% and BSS >7.5 hrs./day and while during dough stage of the crop T min (15-18°C), T max 30°C, RHm 82-85% and BSS more than 8 hrs./day were found conducive for better yield. Cold weather during vegetative phase and warm weather during maturity is desirable for wheat. Long periods of heat stress (above 35°C) conditions in wheat crop during crown root initiation, flowering, and grain filling stages can cause significant yield reduction and may lead to total crop damage. Similar responses to high night temperature have been found under field conditions from stem elongation

to anthesis [9] and from post-anthesis to maturity [10] (Fig. 2).

3.3 Congenial Weather Requirements Infestation of Insect and Diseases of Wheat at Sultanpur

T max 20-29°C and T min 14-8°C, RHm 90-94% and RHe 49-65% and rainfall of 1.5- 11 mm from met. no. week 46th to 4th met. week and T max 20-28.5°C, T min 12-17°C, RHm 88-91% and RHe 42-52% from 1st to 10th met. week at booting and milking stage, were found conducive weather conditions for army worm with no of larva/plant ranging from 0.51-0.57% (Table 3) and leaf blight with disease intensity ranging from 9.8%-34.5% (Table 4), respectively, in wheat crop at district Sultanpur (Fig. 2). It is clear from the Table 4 that 9.8% disease intensity of the leaf blight was found in met. Week 1st with max temperature 19.2°C while the intensity reduced to 6.45% with decrease in max temperature (18.7°C). The table also denotes that as the temperature increases with increase in rainfall from 3rd met week to 10th met week the disease intensity of leaf blight also increases. Similar results are also concluded by Devi et al. [11].

Table 3. Weather parameters for army worm of wheat (Averaged over 5 years)

Std.MET. Week No.	T max(°C)	T min(°C)	RHm %	RHe %	RF (mm)	No. of larva/plant %
46	29.1	14.0	90.9	49.5	1.5	0.51
47	27.5	11.2	88.8	43.9	0.0	0.52
48	28.8	12.3	90.5	49.9	0.0	0.54
49	24.0	9.9	93.5	54.2	0.0	0.59
50	23.1	9.4	88.1	54.5	3.7	0.3
51	21.8	6.6	92.1	57.8	0.0	0.32
52	20.8	6.5	94.0	63.2	0.0	0.38
1	19.2	7.9	93.2	65.4	1.8	0.4
2	18.7	6.3	91.9	62.1	1.1	0.46
3	20.0	6.6	93.9	60.4	3.5	0.52
4	21.6	7.9	91.8	59.2	11.3	0.57

Table 4. Weather parameters for disease intensity of leaf blight (Averaged over 5years)

Std. MET. Week No.	T max(°C)	T min(°C)	RHm %	RHe %	RF (mm)	Leaf blight Diseaseintensity %
1	19.2	7.9	93.2	65.4	1.8	9.8
2	18.7	6.3	91.9	62.1	1.1	6.45
3	20.0	6.6	93.9	60.4	3.5	19.35
4	21.6	7.9	91.8	59.2	11.3	17.1
5	22.7	7.9	90.9	54.6	0.5	20.25
6	23.4	7.9	91.2	50.4	1.5	5.9
7	24.7	9.5	87.7	50.4	0.5	6.3
8	27.4	12.0	87.7	47.2	9.2	20.6
9	27.2	12.9	88.5	47.8	8.3	40.7
10	28.8	12.6	85.1	49.1	10.2	34.5




Fig. 2. Crop weather calendar for wheat

4. CONCLUSION

Crop weather calendar of wheat crop contains a detailed information on wheat crop regarding its climatic requirements from sowing of crop to till harvesting of crop with additional information about the favorable weather conditions of major attacking pest and disease of wheat viz. army worm and leaf blight. Crop contingency plans and Agromet advisory services are variety of the measures to tackle this sort of situations which requires understanding of the crop phenology and effect of weather parameters on crop growth. Crop weather calendar assumes great importance in this scenario. The crop weather calendars visiting be useful to crop insurance personnel in identifying critical stages and appropriate weather indices. It will be of immense use in preparing Agromet advisory bulletins, crop contingency plans and development of insurance products.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

- Porter JR, Gawith M. Temperatures and the growth and development of wheat: A review. *Eur. J Agron.* 1998;10:23–26.
- Prasad PVV, Pisipati SR, Ristic Z, Bukovnik U, Fritz AK. Impact of nighttime temperature on physiology and growth of spring wheat. *Crop Sci.* 2008;48:2372–2380.
- Varshneya MC, Pillai PB. Text book of agricultural meteorology. ICAR. 2008;184–185.
- Kaur P, Bala A, Singh H, Sandhu SS. Guidelines to prepare crop weather calendar. All India Coordinated Research Projection Agrometeorology. School of climate change and agricultural meteorology. Punjab Agricultural University, Ludhiana. 2013;18.
- Praveen KM, Mehera B, Naik J, Gautam S, Madhu BM. Agrometeorological Indices Requirement of Wheat Crop at Allahabad Region under Different Sowing Environment. *Int. J. Curr. Microbiol. App. Sci.* 2018;7(09):2986-2992.
- Wheeler TR, Hong TD, Ellis RH, Batts GR, Morison JIL, Hadley P. The duration and rate of grain growth, and harvest index, of wheat (*Triticum aestivum* L.) in the impact of extreme heat and frost events on wheat crop production: a review. response to temperature and CO₂. *J. Exp. Bot.* 1996;47:623–630.
- Rao VUM, Rao S, AVM, Chandran SMA, Kaur P, Kumar PV, Rao BB, Khandgonda IR, Rao CS. District level crop weather calendars of major crops in India. All India Coordinated Research Project on Agrometeorology, ICAR-Central Research Institute for Dryland Agriculture, Santoshnagar, Hyderabad – 500059; 2015.
- Narayanan S, Prasad PVV, Fritz AK, Boyle DL, Gill BS. Impact of high night-time and high daytime temperature stress on winter wheat. *J Agron Crop Sci.* 2015;201:206–218.
- García GA, Dreccer MF, Miralles DJ, Serrago RA. (High night temperatures during grain number determination reduce wheat and barley grain yield: a field study. *Glob Chang Biol.* 2015;21:4153–4164.
- García GA, Serrago RA, Dreccer MF, Miralles DJ. Post-anthesis warm nights reduce grain weight in field-grown wheat and barley. *Field Crops Res.* 2016;195:50–59.
- Devi HM, Mahapatra S, Dutta S, Das S. Influence of phenological growth stages and meteorological parameters on leaf blight infestation of Wheat in Gangetic plains of West Bengal. *J. Wheat Research.* 2017; 9(2):101-107.

© 2022 Pandey et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:

<https://www.sdiarticle5.com/review-history/82526>