



# Optimizing Nutrient Management for Pearl Millet: Insights from a Long-term Field Experiment

Indra Raj Yadav <sup>a\*</sup>, S. K. Trivedi <sup>a</sup>, Shashi S. Yadav <sup>a</sup>,  
Suwa Lal Yadav <sup>b</sup>, P. A. Khambalkar <sup>a</sup>, Babu Lal Garhwal <sup>a</sup>  
and Anamika Tomar <sup>a</sup>

<sup>a</sup> Department of Soil Science, College of Agriculture, Rajmata Vijayaraje Sciendia Krishi Vishwa Vidyalyaya, Gwalior (Madhya Pradesh)-474 002, India.

<sup>b</sup> School of Agricultural Sciences, Raffles University, Neemrana, Alwar, Rajasthan-301 020, India.

## Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

## Article Information

DOI: <https://doi.org/10.9734/ajsspn/2024/v10i3347>

## Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/121335>

Original Research Article

Received: 30/05/2024

Accepted: 01/08/2024

Published: 05/08/2024

## ABSTRACT

The study investigates the impact of integrated nutrient management strategies on the yield and nutrient uptake of pearl millet (*Pennisetum glaucum*), a vital crop in arid and semi-arid regions. Pearl millet is a significant source of dietary energy and essential nutrients for rural populations in India. However, its productivity remains low compared to other millet-producing countries. This research aims to enhance pearl millet yield through a combination of organic manures, biofertilizers, and chemical fertilizers, addressing the need for sustainable agricultural practices. The experiment was conducted at the Research Farm of Rajmata Vijayaraje Scindia Krishi Vishwavidyalaya during

\*Corresponding author: Email: [indrarajyadav100@gmail.com](mailto:indrarajyadav100@gmail.com);

**Cite as:** Yadav, Indra Raj, S. K. Trivedi, Shashi S. Yadav, Suwa Lal Yadav, P. A. Khambalkar, Babu Lal Garhwal, and Anamika Tomar. 2024. "Optimizing Nutrient Management for Pearl Millet: Insights from a Long-Term Field Experiment". *Asian Journal of Soil Science and Plant Nutrition* 10 (3):369-75. <https://doi.org/10.9734/ajsspn/2024/v10i3347>.

the kharif seasons of 2022 and 2023. Twelve treatment combinations were tested, including 100% NPK along with farmyard manure (FYM) and seed treatments with biofertilizers. Results indicated that the treatment combining 100% NPK with FYM at 10 t/ha/year and biofertilizers significantly improved biological yield, nutrient uptake, and soil organic carbon content. Specifically, the highest biological yield recorded was 8026 kg/ha, with notable increases in total nitrogen, phosphorus, and potassium uptake. This study underscores the importance of integrated nutrient management in enhancing pearl millet productivity while maintaining soil health. By adopting these practices, farmers can improve crop yields and contribute to food security in regions reliant on pearl millet. The findings provide a framework for future research and practical applications in sustainable agriculture, emphasizing the need for balanced nutrient supply systems to optimize crop performance and soil fertility.

**Keywords:** *Azotobacter*; FYM; long-term; physico-chemical properties.

## 1. INTRODUCTION

Pearl millet (*Pennisetum glaucum*) is commonly known as Bajra, Bajri, Sajje, Kambu, Kamban, Sajjalu, etc. in various Indian local languages and is commonly used for food, feed, and forage. It is the most widely grown millet and it is one of the important millet crops in hot and dry areas of arid and semi-arid climatic conditions, particularly in Rajasthan. Pearl millet belongs to the family Gramineae or Poaceae. The inflorescence of pearl millet is known as spike. It is a highly cross-pollinated crop due to protogynous habit and C<sub>4</sub> plant, it has very high photosynthetic efficiency and dry matter production capacity. Pearl millet originated in tropical Western Africa about 4000 years ago. India is the largest producer of pearl millet, among all the states and UTs, Rajasthan (46%), Maharashtra (19%), Gujarat (11%), Uttar Pradesh (8%) and Haryana (6%) [1] was among top pearl millet producing states.

In India, pearl millet is a primary source of dietary energy (360 kcal/kg) for rural populations and the fourth most important cereal after rice, wheat, and sorghum [2] It is a rich source of protein, calcium, phosphorous, and iron. Pearl millet grain contains fairly high amounts of thiamine, riboflavin, and niacin. Pearl millet grain is also used for non-food purposes such as poultry feed, cattle feed, and alcohol extraction [3]. The productivity of pearl millet is very low compared to important pearl millet-growing countries in the world. It is, therefore, necessary to increase the production and productivity of pearl millet by adopting scientific innovations [4].

An integrated nutrient supply system comprising biofertilizers, organic manures and FYM in conjunction with chemical fertilizers is necessary to meet the crop nutrient demand. In the research problem, various organic like FYM and

seed treatment with PSB and Rhizobium and inorganic nutrient sources like nitrogen, phosphorus, potassium, sulphur, zinc and ferrous containing fertilizer were used. Long-term application of inorganic fertilizers depletes the physical, chemical, and biological qualities of soil and pollutes the environment when combined with organic additions [5]. In addition to providing the soil with nutrients and organic matter, organic manures also influence the soil's structure, turnover of nutrients, biodiversity, and activity of the microbial population, among many other changes in the physical, chemical, and biological characteristics of the soil. The physico-chemical qualities of soils can be enhanced by the use of organic manures alone or in conjunction with chemical fertilizers [6]. A judicious combination of chemical fertilizers, organic manures and bio-fertilizers should be formulated for crops and cropping system within the ecological, social and economic possibilities [7].

Since information pertaining to the above aspects is meagre, the present investigation was carried out to study the effect of integrated nutrient management (INM) on nutrient uptake, productivity and soil sustainability to enhance the fertility and productivity of the soil.

## 2. MATERIALS AND METHODS

The present investigation is a part of an ongoing long-term field experiment of integrated nutrient management on pearl millet. The experiment was initiated in 2002 on permanent plots with latitude of 26° 13'N and longitude 76° 10'E with an altitude 197 meters. The climate of the experimental site is semi-arid and subtropical with extreme weather conditions having hot and dry summer and cold winter, where maximum temperature goes up to 45 °C during summer and minimum

as low as 2.80 °C. The mean annual rainfall of the area is 700-800 mm. The soil of the experimental field is alluvial, sandy clay loam in texture and classified as Typic Ustochrepts at great group level comes under Inceptisols. The experiment was laid out with randomized block design having three replications comprising of 12 treatments, viz T<sub>1</sub>-Absolute Control, T<sub>2</sub>-100% N, T<sub>3</sub>-100% NP, T<sub>4</sub>-100% NPK, T<sub>5</sub>-100% NPK + 25 kg ZnSO<sub>4</sub>/ha/yr, T<sub>6</sub>-100% NPK + 50 Kg FeSO<sub>4</sub>/ha/yr, T<sub>7</sub>-100% NPK + FeSO<sub>4</sub> + ZnSO<sub>4</sub>, T<sub>8</sub>-100% NPK + 1% (FeSO<sub>4</sub> + ZnSO<sub>4</sub> Spray), T<sub>9</sub>-50% NPK + FYM @ 10t/ha/yr, T<sub>10</sub>-75% NPK + FYM @ 10t/ha/yr, T<sub>11</sub>-100% NPK+ FYM @ 10t/ha/yr and T<sub>12</sub>-100% NPK +FYM @ 10 t/ha/yr + {PSB+ Azotobacter (Seed treatment)}, respectively.

The recommended dose of fertilizer for pearl millet was 80 N + 40 P<sub>2</sub> O<sub>5</sub> + 20 K<sub>2</sub>O kg/ha. Half of the nitrogen was applied in the form of urea as a basal dose and the remaining was top-dressed after 1<sup>st</sup> irrigation at 30 DAS. Full dose of phosphorus and potash applied as basal through single super phosphate and muriate of potash. The treatments were applied as per treatment details and FYM was applied 1 month before sowing. The azotobacter was applied as seed treatment at the rate of 10g/ kg seed. The JBV-3, cultivar was sown 5 kg/ha seed rate during July month and harvested in the second week of October. The pH, EC and OC content in soil before sowing were 7.72, 0.23 dSm<sup>-1</sup>, and 0.39 % respectively.

### 3. RESULTS AND DISCUSSION

The data furnished (Table1) that the test weight and harvest index were found non-significant in both years. The data showed that the pooled mean found significantly higher biological yield (Table 1) (8026 kg/ha) were recorded with treatment T<sub>12</sub>-100% NPK +FYM @ 10 t/ha/yr + {PSB+ Azotobacter (Seed treatment), closely followed by T<sub>11</sub>-100% NPK+ FYM @ 10t/ha/yr and T<sub>10</sub>-75% NPK + FYM @ 10t/ha/yr (7380 and 6702 kg/ha). The minimum biological yield was recorded with the control treatment (2987 kg/ha). The application of the organic and inorganic nutrient sources including 100% NPK + FYM + PSB+ Azotobacter (Seed treatment) provided balanced nutrients to the pearl millet which resulted into significant enhancement in biological yield. Similar result also reported by Abraham et al. [8], Devi et al. [9], Ram and Dhaliwal [10], Kumar et al. [11] and Susmitha et al. [12].

The data obtained that total nitrogen uptake (Table 2) significantly higher the pooled mean were recorded with T<sub>12</sub>-100% NPK +FYM @ 10 t/ha/yr + {PSB+ Azotobacter (Seed treatment) (89.84 kg/ha), closely followed by T<sub>11</sub>-100% NPK+ FYM @ 10t/ha/yr and T<sub>10</sub>-75% NPK + FYM @ 10t/ha/yr (78.39 and 67.41 kg/ha). The minimum total nitrogen uptake was recorded with the control treatment (20.87 kg/ha). This might due to the addition of FYM enhanced significantly total nitrogen uptake by the crop. More uptake of N under the treatment with FYM resulted in mineralized N from FYM could meet the need of nutrients of crop. The outcome is turned with findings by Dahiya et al. [13] and Singh et al. [14]. The data obtained that pooled mean of total phosphorus uptake (Table 2) maximum recorded with T<sub>12</sub>-100% NPK +FYM @ 10 t/ha/yr + {PSB+ Azotobacter (Seed treatment) (17.90 kg/ha), closely followed by T<sub>11</sub>-100% NPK+ FYM @ 10t/ha/yr and T<sub>10</sub>-75% NPK + FYM @ 10t/ha/yr (15.97 and 13.25 kg/ha). The minimum total phosphorus uptake was recorded with control treatment (4.05 kg/ha). This might be due to the higher availability of P from applied FYM resulting in the decomposing of the organic materials which produced from the solubilizing action of organic acid. The data obtained that pooled mean of total potassium uptake (Table 2) maximum recorded with T<sub>12</sub>-100% NPK +FYM @ 10 t/ha/yr + {PSB+ Azotobacter (Seed treatment) (111.50 kg/ha), closely followed by T<sub>11</sub>-100% NPK+ FYM @ 10t/ha/yr and T<sub>10</sub>-75% NPK + FYM @ 10t/ha/yr (100.07 and 87.40 kg/ha). The minimum total potassium uptake was recorded with control treatment (30.27 kg/ha). These results also confirmed by Ramdas et al. [15], Choudhary et al. [16], Yadav et al. [17] and Sunag et al. [18].

The data furnished (Table 3) that the pH and electrical conductivity were found to be non-significant in both years. The data presented that the pooled mean found significantly higher (Table 3) organic carbon recorded with T<sub>12</sub>-100% NPK +FYM @ 10 t/ha/yr + {PSB+ Azotobacter (Seed treatment) (0.408 %), it was at par with T<sub>11</sub>-100% NPK+ FYM @ 10t/ha/yr (0.405 %). The minimum organic carbon recorded with control treatment with (0.373 %). The organic carbon content of the soil due to the application of Farm Yard Manure might be due to the deposition of organic matter in the soil in the crop residues and the direct addition of OM through FYM along with inorganic fertilizer. Similar concluded by Kumar et al. [19], Rao et al. [20], Bairwa et al. [21] and Kalaliya et al. [22].

**Table 1. Effect of integrated nutrient management on yield attributes and yield of pearl millet**

Treatments	Test weight (g)			Harvest index (%)			Biological yield (kg/ha)		
	2022	2023	Pooled	2022	2023	Pooled	2022	2023	Pooled
T <sub>1</sub>	7.46	7.45	7.46	30.69	29.78	30.24	2917	3056	2987
T <sub>2</sub>	7.52	7.50	7.51	31.22	30.71	30.96	3574	3623	3599
T <sub>3</sub>	7.54	7.53	7.54	31.27	30.58	30.92	4431	4575	4503
T <sub>4</sub>	7.58	7.62	7.60	28.65	28.45	28.55	5324	5486	5405
T <sub>5</sub>	7.62	7.61	7.62	28.11	28.54	28.32	5534	5496	5515
T <sub>6</sub>	7.60	7.60	7.60	28.58	28.27	28.43	5353	5466	5409
T <sub>7</sub>	7.63	7.62	7.63	28.78	28.76	28.77	5596	5637	5617
T <sub>8</sub>	7.61	7.60	7.61	28.42	28.40	28.41	5189	5478	5334
T <sub>9</sub>	7.65	7.66	7.66	27.78	27.53	27.65	6049	6093	6071
T <sub>10</sub>	7.69	7.72	7.71	28.26	27.57	27.91	6549	6856	6702
T <sub>11</sub>	7.50	7.52	7.51	29.14	29.64	29.39	7362	7398	7380
T <sub>12</sub>	7.50	7.50	7.50	29.45	29.27	29.36	7988	8064	8026
S. Em. (±)	0.319	0.343	0.234	1.275	1.192	0.873	213.86	245.46	162.78
CD (0.05%)	NS	NS	NS	NS	NS	NS	627.24	719.93	463.95

**Table 2. Effect of integrated nutrient management on total uptake by pearl millet**

Treatments	Total nitrogen uptake (kg/ha)			Total phosphorus uptake (kg/ha)			Total potassium uptake (kg/ha)		
	2022	2023	Pooled	2022	2023	Pooled	2022	2023	Pooled
T <sub>1</sub>	20.47	21.26	20.87	3.34	4.77	4.05	30.52	30.02	30.27
T <sub>2</sub>	30.92	31.34	31.13	4.74	6.40	5.57	39.15	39.74	39.45
T <sub>3</sub>	38.83	39.97	39.40	7.60	9.98	8.79	48.96	50.71	49.84
T <sub>4</sub>	47.99	48.86	48.43	8.74	12.13	10.44	66.06	67.56	66.81
T <sub>5</sub>	49.62	49.61	49.62	8.87	12.24	10.56	69.22	68.45	68.84
T <sub>6</sub>	49.74	50.21	49.98	9.00	12.30	10.65	66.50	68.24	67.37
T <sub>7</sub>	52.14	53.67	52.91	9.57	13.14	11.36	70.34	70.87	70.61
T <sub>8</sub>	49.58	49.95	49.77	8.99	12.35	10.67	68.47	76.47	72.47
T <sub>9</sub>	57.31	57.86	57.58	9.24	12.94	11.09	74.69	78.06	76.37
T <sub>10</sub>	66.13	68.69	67.41	11.12	15.38	13.25	85.02	89.77	87.40
T <sub>11</sub>	78.05	78.73	78.39	13.41	18.52	15.97	99.92	100.21	100.07
T <sub>12</sub>	89.53	90.14	89.84	15.13	20.66	17.90	110.68	112.31	111.50
S. Em. (±)	1.52	1.73	1.15	0.23	0.42	0.24	1.60	1.78	1.20
CD (0.05%)	4.46	5.09	3.29	0.69	1.23	0.68	4.69	5.21	3.41

**Table 3. Effect of integrated nutrient management on pH, Electrical conductivity and Organic carbon in soil after harvest of pearl millet**

Treatments	Soil pH			Electrical conductivity (dSm <sup>-1</sup> )			Organic carbon (%)		
	2022	2023	Pooled	2022	2023	Pooled	2022	2023	Pooled
T <sub>1</sub>	7.70	7.70	7.70	0.119	0.117	0.118	0.372	0.373	0.373
T <sub>2</sub>	7.60	7.60	7.60	0.119	0.118	0.119	0.375	0.378	0.376
T <sub>3</sub>	7.60	7.60	7.60	0.121	0.120	0.121	0.378	0.376	0.377
T <sub>4</sub>	7.60	7.60	7.60	0.123	0.123	0.123	0.389	0.390	0.390
T <sub>5</sub>	7.60	7.60	7.60	0.125	0.125	0.125	0.390	0.391	0.390
T <sub>6</sub>	7.60	7.60	7.60	0.125	0.126	0.126	0.389	0.390	0.390
T <sub>7</sub>	7.60	7.60	7.60	0.126	0.126	0.126	0.391	0.392	0.392
T <sub>8</sub>	7.60	7.60	7.60	0.123	0.123	0.123	0.388	0.389	0.389
T <sub>9</sub>	7.60	7.50	7.55	0.126	0.125	0.126	0.392	0.394	0.393
T <sub>10</sub>	7.50	7.50	7.50	0.125	0.125	0.125	0.395	0.397	0.396
T <sub>11</sub>	7.50	7.50	7.50	0.124	0.124	0.124	0.405	0.404	0.405
T <sub>12</sub>	7.50	7.50	7.50	0.124	0.123	0.124	0.407	0.409	0.408
S. Em. (±)	0.212	0.204	0.147	0.003	0.003	0.002	0.005	0.005	0.003
CD (0.05%)	NS	NS	NS	NS	NS	NS	0.014	0.014	0.009

#### 4. CONCLUSION

On the basis of these findings, it was concluded that treatment T<sub>12</sub>-100% NPK +FYM @ 10 t/ha/yr + {PSB+ Azotobacter (Seed treatment) found best among remaining treatment. This research showed the highest yield was obtained and a significantly impact on soil properties as prospective to soil fertility.

#### DISCLAIMER (ARTIFICIAL INTELLIGENCE)

Author(s) hereby declare that NO generative AI technologies such as Large Language Models (ChatGPT, COPILOT, etc) and text-to-image generators have been used during writing or editing of manuscripts.

#### COMPETING INTERESTS

Authors have declared that no competing interests exist.

#### REFERENCES

1. Amarghade and Singh R. () Effect of inorganic and organic nutrient sources on growth and yield of pearl millet (*Pennisetum glaucum* L.). The Pharma Innovation Journal. 2021;10(10):507-509.
2. Tomar PS, Verma SK, Gupta N, Bansal KN. Effect of long-term integrated nutrient management on productivity of pearl millet (*Pennisetum glaucum*)-mustard (*Brassica juncea*) cropping system and soil fertility in an Inceptisol. Journal of the Indian Society of Soil Science. 2018;66(3):295-299.
3. Mohan N, Singh S. Effect of organic manures and sources of sulphur on growth and yield of kharif pearl millet (*Pennisetum glaucum* L.). The Pharma Innovation Journal. 2022;11(3):2257-61.
4. Thumar CM, Dhdhat MS, Chaudhari NN, Hadiya NJ, Ahir NB. Growth, yield attributes, yield and economics of summer pearl millet (*Pennisetum glaucum* L.) as influenced by integrated nutrient management. International Journal of Agriculture Sciences; 2016. ISSN, 0975-3710.
5. Singh R, Gupta AK, Ram T, Choudhary GL, Sheoran AC. Effect of integrated nitrogen management on transplanted pearl millet (*Pennisetum glaucum*) under rainfed condition. Indian Journal of Agronomy. 2013;58(1):81-85.
6. Prasad J, Karmakar S, Kumar R, Mishra B. Influence of integrated nutrient management on yield and soil properties in maize-wheat cropping system in an Alfisol of Jharkhand. Journal of the Indian Society of Soil Science. 2010;58(2):200-204.
7. Arya V, Trivedi SK, Tomar PS, Singh M, Dhakad H. Effect of Integrated Nutrient Management (INM) on physico-chemical Properties of Soils under Pearl Millet-mustard Cropping Sequence in Typic Ustochrepts. International Journal of Plant & Soil Science. 2022 34(19), 183-193.
8. Abraham T, Thenua OVS, Singh SP, Jacob P. Performance of groundnut as influenced by organic and inorganic sources of nutrients and their method of application. Legume Research. 2008;31: 224-226.
9. Devi KN, Singh MS, Singh NG, Athokpam S. Effect of integrated nutrient management on growth and yield of wheat (*Triticum aestivum* L.). Journal of Crop and Weed. 2011;7(2): 23-27.
10. Ram Hari, Dhaliwal SS. Effect of varieties and integrated nutrient management techniques on growth, productivity, quality and economics of barley (*Hordeum vulgare* L.) International Journal of Agricultural Sciences. 2012;8(1):91-97.
11. Kumar Pradeep, Singh Rakesh, Singh Archana, Paliwal Dinesh, Kumar Sushil) Integrated nutrient management in pearl millet (*Pennisetum glaucum*) – wheat (*Triticum aestivum*) cropping sequence in semi arid condition of India. Internat. J. agric. Sci. 2014;10(1):96-101.
12. Susmitha Kali C Umesha, Shahazad Ahmed Khan. Effect of Biofertilizer and Organic Manure on Growth and Yield of Pearl Millet (*Pennisetum Glaucum* L.)". International Journal of Environment and Climate Change. 2022 12(10):425-31. Available: <https://doi.org/10.9734/ijecc/2022/v12i1030815>.
13. Dahiya SS, Goyal S, Antil RS, Karwasra SPS. Effect of farm yard manure and cadmium on dry matter yield and nutrient uptake by maize. J. Ind. Soc. Soil. Sci. 1987;35:460-464.
14. Singh Vinay, Kumar R, Ram Lakhan. Effect of applied farmyard manure and molybdenum on yield and nutrient uptake by Egyptian clover. Ind. J. Agron. 1994;39(3):307-309.

15. Ramdas MG, Manjunath BL, Singh NP, Ramesh R, Verma RR, Latore AM, Ruenna D'Souza, Barnes N, Kulkarni R. Effect of organic and inorganic sources of nutrients on soil microbial activity and soil organic carbon build-up under rice in west coast of India. *Archivers of Agronomy and Soil Science*; 2016.
16. Choudhary K, Sharma SR, Jat R, Didal VK. Effect of organic manures and mineral nutrients on quality parameters and economics of sesame (*Sesamum indicum* L.). *Journal of Pharmacognosy and Phytochemistry*. 2017;6(3):263-265.
17. Yadav, L.C., Yadav L.R., Yadav Nagesh and Yadav Mamta. (2018). Effect of organic sources of nitrogen fertilization on yield and yield attributes pearl millet [*Pennisetum glaucum* (L.) R. BR. emend stuntz]. *International Journal of Chemical Studies*. 6(3): 2329-2331.
18. Sunag MN, Singh RM, Singh V, Singh K. Effect of integrated nutrient management on the performance of barley (*Hordeum vulgare* L.). *The Pharma Innovation Journal*. 2021;10(9):1492-1496.
19. Kumar S, Dahiya R, Kumar P, Jhorar BS, Phogat VK. Long Term effect of organic materials and fertilizers on soil properties in pearl millet - wheat cropping system. *Ind. J. Agric. Sci*. 2012 46(2): 161-166.
20. Rao PM, Sagar V, Suresh K, Padmaja G, Reddy SN. Influence of organic and inorganic sources of nitrogen on nutrient uptake and yield of maize in maize-mustard cropping system in Northren Telangana zone. *The Pharma Innovation Journal*. 2021;10(10):1351-1357.
21. Bairwa J, Dwivedi BS, Rawat A, Thakur RK, Mahawar Neeta. Long-Term Effect of Nutrient Management on Soil Microbial Properties and Nitrogen Fixation in a Vertisol under Soybean- Wheat Cropping Sequence. *Journal of the Indian Society of Soil Science*. 2021;69(2):171-178.
22. Kalaliya A, Sharma SK, Kamboj BR. Effect of nutrient management on growth, yield attributing characters, yield and economics of rainfed pearl millet [*Pennisetum glaucum* (L) R.Br.]. *The Pharma Innovation Journal*. 2022;11(7):290-294.

**Disclaimer/Publisher's Note:** The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of the publisher and/or the editor(s). This publisher and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.

© Copyright (2024): Author(s). The licensee is the journal publisher. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:

<https://www.sdiarticle5.com/review-history/121335>